

Opinion

Common Pathogenetic Mechanisms Underlying Aging and Tumor and Means of Interventions

Weiye Shen^{1,2,3}, Jiamin He^{1,2,3}, Tongyao Hou^{1,2,3*}, Jianmin Si^{1,2,3*}, Shujie Chen^{1,2,3*}

¹Department of Gastroenterology, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou 310016, Zhejiang, China. ²Institute of Gastroenterology, Zhejiang University, Hangzhou 310016, Zhejiang, China. ³Prevention and Treatment Research Center for Senescent Disease, Zhejiang University School of Medicine, Zhejiang, China

[Received September 9, 2021; Revised December 6, 2021; Accepted December 7, 2021]

ABSTRACT: Recently, there has been an increase in the incidence of malignant tumors among the older population. Moreover, there is an association between aging and cancer. During the process of senescence, the human body suffers from a series of imbalances, which have been shown to further accelerate aging, trigger tumorigenesis, and facilitate cancer progression. Therefore, exploring the junctions of aging and cancer and searching for novel methods to restore the junctions is of great importance to intervene against aging-related cancers. In this review, we have identified the underlying pathogenetic mechanisms of aging-related cancers by comparing alterations in the human body caused by aging and the factors that trigger cancers. We found that the common mechanisms of aging and cancer include cellular senescence, alterations in proteostasis, microbiota disorders (decreased probiotics and increased pernicious bacteria), persistent chronic inflammation, extensive immunosenescence, inordinate energy metabolism, altered material metabolism, endocrine disorders, altered genetic expression, and epigenetic modification. Furthermore, we have proposed that aging and cancer have common means of intervention, including novel uses of common medicine (metformin, resveratrol, and rapamycin), dietary restriction, and artificial microbiota intervention or selectively replenishing scarce metabolites. In addition, we have summarized the research progress of each intervention and revealed their bidirectional effects on cancer progression to compare their reliability and feasibility. Therefore, the study findings provide vital information for advanced research studies on age-related cancers. However, there is a need for further optimization of the described methods and more suitable methods for complicated clinical practices. In conclusion, targeting aging may have potential therapeutic effects on aging-related cancers.

Key words: aging, cancer, similarities, pathogenetic mechanisms, interventions

Aging of the human population is gradually becoming an important global issue. According to the United Nations, the global human population aged >60 years will increase by two-fold by 2050 compared to that in 2000 [1]. Aging of the human population is gradually becoming an important global issue. According to the United Nations, the global human population aged >60 years will increase by two-fold by 2050 compared to that in 2000 [2]. Therefore, aging causes severe stress on the human health

system. Therefore, there is an urgent need to improve the health of this population. To explore the mechanism of aging, several studies have found several common hallmarks of aging in different individuals, such as genomic alteration, epigenetic changes, mitochondrial damage, and cellular senescence. The findings of previous studies have provided clues to further explore the mechanisms of anti-aging or alleviating aging-related damage and cancer [3].

*Correspondence should be addressed to: Dr. Shujie Chen (Email: chenshujie77@zju.edu.cn), Dr. Jianmin Si (Email: jianmin_si@zju.edu.cn) and Dr. Tongyao Hou (Email: houtongyao0805@zju.edu.cn), Department of Gastroenterology, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou 310016, Zhejiang, China. #These authors contributed equally to this work.

Copyright: © 2021 Shen W et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Recently, the high rates of morbidity and mortality caused by various cancers have made it a much-researched topic. According to statistics, the incidence of cancer among humans is 11-fold higher in individuals aged >65 years than in individuals aged < 65 years. Furthermore, most cancers are diagnosed in people aged >55 years [4]. Currently, the gap in the incidence of cancer between young and older individuals has increased compared to that described earlier. In some cancers, aging is a factor that leads to malignant tumors with a poorer prognosis; hence, various cancers seem to have a close relationship with aging [5, 6].

A recent study performed by Gomes et al. revealed that aging-mediated metabolic reconstitution, especially accumulation of methylmalonic acid, can form an internal environment that facilitates cancer cell growth and progression [7]. These findings further confirmed the existence of a mysterious intersection between aging and cancer. Therefore, exploring the pathogenesis of aging-induced cancer is of great importance. In this study, the junctions of aging and cancer were addressed by comparing various research findings. The study results indicate that delaying aging may be a promising intervention against aging-related cancers.

Aging and cancer have common pathogenetic mechanisms

The mechanisms of the occurrence and development of aging and cancer are being researched since the past few decades, and several results have been achieved. These studies have revealed several mechanisms involved in the occurrence and/or development of both aging and cancer. For instance, studies have reported that cellular senescence significantly diminishes in malignant tumors, enhanced ubiquitin–proteasome and lysosome–autophagy systems prolong lifespan but promote cancer progression, and alterations in the systemic immune microenvironment mediated by aging push further demic recession and cancer appearance. In addition, intestinal flora, local or systemic metabolism, and hormones change with age, which in turn affects the longevity of human, drives tumorigenesis, and causes cancer deterioration in health simultaneously. Moreover, aging and cancer are regulated by the expression of similar genes. Therefore, we aimed to precisely describe the common mechanisms involved in the occurrence and development of both aging and cancer (Fig. 1, Table 1, and Table 2).




Figure 1. Common pathogenetic mechanisms between aging and cancer.

Cellular senescence: What happens in cancer?

Cellular senescence refers to the permanent proliferative arrest of cells subjected to stressors. It can be induced by various physiological and pathological factors, such as telomere erosion, oxidative stress, and activation of oncogenes [8].

In cancer cells, powerful telomerase activity leads to attenuated telomere erosion and delayed cellular

senescence. Therefore, antitelomerase therapy can be an important therapeutic intervention for malignant growth and can induce cell death. Notably, cancers that retain wild-type *p53*, a well-known anti-oncogene, can falsify the response to antitelomerase therapies, which only trigger proliferative arrest rather than inducing cell death. Furthermore, this change results in prolonged treatment and increased drug resistance [9].

Table 1. Common mechanism between aging and cancer.

Mechanism	Alteration during aging	Effect on aging	Effect on cancer
Proteostasis			
Chaperones	Decrease	Delay	Promote
Ubiquitin-proteasome	Decrease	Delay	Promote
Lysosome-autophagy	Decrease	Delay	Promote
Microbiota disorder			
Akkermansia	Decrease	Delay	Inhibit
Bifidobacterium	Decrease	Delay	Inhibit
Escherichia coli	Increase	/	Promote
Bacteroides fragilis	Increase	/	Promote
Immune microenvironment			
Inflammation	Increase	Accelerate	Promote
Immunosenescence	Increase	/	Promote
Metabolic disorder			
Energy metabolism			
NAD+	Decrease	Delay	Promote
Mitochondrial damage	Increase	Accelerate	Promote
Substance metabolism			
Methylmalonic acid	Increase	/	Promote
Quinolate	Increase	/	Promote
Phosphoenolpyruvate	Increase	/	Inhibit
α -ketoglutarate	Decrease	Delay	Bidirectional
Endocrine disorder			
TSH	Decrease	Delay	Promote
GH/IGF-1	Decrease	Accelerate	Promote
FSH/LH	Decrease	Delay	Promote
Genetics			
Klotho	Low expression	Delay	Inhibit
AUF1	/	Delay	Bidirectional
SIRT1	Low expression	Delay	Bidirectional
P16	High expression	Accelerate	Inhibit

NAD+, nicotinamide adenine dinucleotide; TSH, thyroid stimulating hormone; GH, growth hormone; IGF-1, insulin-like growth factor I; FSH, follicle-stimulating hormone; LH, luteinizing hormone.

The life and death of cells is largely determined by their redox status. Oxidative stress is a cause of cellular senescence in normal cells. In malignant cells, although excessive proliferation results in higher generation of reactive oxygen species (ROS), the cells can realign their

redox status and enhance their antioxidant ability, thereby avoiding reaching the ROS thresholds that trigger cellular senescence. Moreover, the cells can also optimize ROS-mediated DNA damage and subsequent malignant proliferation [10].

Table 2. Common phenomena between aging and cancer.

Phenomenon	Alteration during aging	Alteration in cancer
Cellular senescence	Increase	Increase (pre-malignant tumor) Decrease (malignant tumors)
Epigenetics		
DNA Methylation age	Accelerate	Accelerate
Genomic imprinting	Increase/Decrease	Increase/Decrease
Gene silencing	Loss	Loss

In pre-malignant tumors, activation of oncogenes, such as *HRASG12V* and *EGFR*, leads to permanent cell proliferative arrest and inhibits cancer progression [11, 12]. In other words, cellular senescence serves as a protective mechanism and extensively exists in the precancerous stage. However, during the stage of tumor

progression and transformation to malignancy, inactivation of anti-oncogenes and common overexpression of other oncogenes disables the process of cellular senescence [13] (Fig. 2).




Figure 2. Disabling cellular senescence in cancer. 1. Powerful telomerase activity attenuates telomere erosion; 2. Realigning redox improves antioxidant capacity; 3. Although activation of individual oncogenes triggers cell proliferative arrest in the early stage, subsequent inactivation of anti-oncogenes and activation of multiple oncogenes restores cell proliferation.

Alterations in proteostasis links aging with cancer

Chaperones, ubiquitin–proteasome, and lysosome–autophagy systems are important pillars of proteostasis. ATP-dependent chaperones are significantly inhibited during the aging process because of damage to energy metabolism. A research study on the nervous system revealed that 32% ATP-dependent chaperones were downregulated in the aged brain [14]. In addition, there was also a sharp decrease in the chaperones that play a role in target recognition [15]. In older individuals, a marked decline in autophagic function has also been shown in various species [16, 17], which may be due to a decline in the expression of *Atg* and *Sirtuin 1* or hyperactivation of TORC1 [18]. Moreover, repression of proteasome activity has been observed in the early stages of various organic aging processes, such as aging of the brain and skin [19, 20].

In contrast, alterations in proteostasis also influence the process of aging. In yeast, increasing proteasome capacity by upregulating the transcription factor Rpn4 can prolong their replicative lifespan [21]. In *Drosophila melanogaster*, upregulation of Rpn11 and DmPI31, gamma irradiation, and other processes have been shown to extend the lifespan by activating the proteasome [22–24]. In mice, the phenomenon of delaying aging by upregulation of IGF-1 or dietary intervention has been shown to be closely related to proteasome activation [25, 26]. In humans, powerful proteasome activity has also been observed in centenarians [27]. Similarly, most longevity methods have been shown to enhance

autophagic function [28]. According to Krøll et al., RNA chaperones are involved in cell immortalization [29]. Furthermore, Ito et al. reported that improving mitochondrial chaperones could lead to longevity in elegans [30].

However, although enhanced proteasome capacity and autophagic function can extend the lifespan, it seems to promote cancer progression. For instance, the proteasome activator REGγ can enhance the transforming growth factor (TGF)-β pathway and result in lung cancer metastasis [31]. Tribble homolog 2 promotes ubiquitin degradation and weakens oxidative damage by upregulating proteasome activity, which promotes the progression of liver cancer [32]. Furthermore, *Fusobacterium nucleatum* has been shown to induce chemoresistance by activating autophagy in colorectal cancer (CRC) [33]. Autophagy also triggers immune evasion by improving MHC-I degradation in pancreatic malignant tumors [34]. In addition, cachexia, ubiquitin–proteasome, and lysosome–autophagy systems are significantly activated in gastric cancer [35]. Moreover, heat shock proteins (HSPs), especially HSP90 and HSP70, play an important role in folding and transporting pivotal cancer proteins and are indispensable factors in cancer aggressiveness [36].

In conclusion, it was evident that powerful ubiquitin–proteasome and lysosome–autophagy systems contribute to better proteostasis and longevity, thus inducing the degradation of anti-cancer factors and worsening of cancer.

Roles of microbiota disorders in aging and cancers

The intestinal microbiota plays an indispensable role in the degradation of carbohydrates, synthesis of short-chain fatty acids, amino acids, and vitamins [37]. Maintaining a normal composition and proportion of intestinal microbiota is the foundation of the integrity of the epithelial barrier and immune balance. In the process of human aging, the composition of the gut microbiota changes constantly and manifests as a decline in microbial diversity, with a decrease in *Bifidobacterium* and increase in *Clostridium*, *Enterobacteriaceae*, and *Enterococci* [38]. In aged animals, researchers have observed decreased genera, indicating an antiphlogistic and butyrate-producing action, as well as increased genera, resulting in the degradation of mucin [39]. These changes can be used to explain aggravated gut inflammation and impaired integrity of the gut epithelial barrier in older individuals.

Recently, researchers have revealed the role of intestinal flora in aging. For instance, in a previous study among centenarians and supercentenarians, there was an abundance of *Christensenella*, *Akkermansia*, and *Bifidobacterium* [40]. In addition, researchers have observed an increase in the expression of genes regulating xenobiotic biodegradation and metabolism such as toluene, chlorocyclohexane, and caprolactam, and a reduction in the expression of genes manipulating the carbohydrate metabolism in centenarians and supercentenarians [41]. Therefore, it is evident that certain flora may prolong one's lifespan. A recent study has also proved this hypothesis, showing that administration of *Akkermansia muciniphila* by oral gavage prolonged the lifespan of progeroid mice [42]. Furthermore, *A. muciniphila* supplementation successfully delayed aging-mediated reduction of colonic mucus thickness [43].

In addition, the intestinal flora possibly intervenes in tumorigenesis and disease progression. Previous studies have indicated differences in the intestinal microbiota between healthy people and patients with cancer, which present as a reduction in the number of beneficial bacteria and an increase in the number of opportunistic pathogens in the gut of patients with cancer [44]. In subsequent experiments, most bacteria were proven to have a positive or negative correlation with cancer.

Evidence shows that *Fusobacterium* stimulates the growth and metastasis of CRC and breast cancer [45-47]. In 2020 year, our team further discovered that *Fusobacterium* motivated the migration of CRC cells and lung metastasis by upregulating the levels of the long non-coding RNA Keratin7-antisense and Keratin7 [48]. In addition, the main mechanisms that have been explored are also involved in virulence factors, immunoregulation, microRNAs, and bacterial metabolism [49]. In addition,

colibactin-producing *Escherichia coli* has been shown to be carcinogenic by suppressing effective T-cell response [50]. Moreover, enterotoxic *Bacteroides fragilis* can accelerate tumor growth by enhancing Treg activity or mediating the long non-coding RNA BFAL1, which depends on the RHEB/mTOR pathway [51]. A recent study has also proposed that seropositivity to seven common *E. coli* and two common enterotoxic *B. fragilis* was closely related to CRC progression, especially that of proximal colon cancers [52]. In addition to an increase in the carcinogenic flora, a decrease in probiotics has also been observed in patients with cancer. Therefore, it is evident that there is a close relationship between chronic colitis and CRC. This year, our team revealed that *B. adolescentis* can suppress DSS-induced chronic colitis by enhancing the protective Treg/Th2 response and promoting intestinal flora remodeling [53]. Moreover, based on test data from some references, *Bifidobacteria* can reduce the occurrence and progression of CRC by downregulating EGFR, HER-2, and COX-2 [54], manipulating microRNAs [55, 56], regulating immune response and so on [57]. *Faecalibacterium prausnitzii* can also inhibit the development of breast cancer by downregulating the interleukin (IL)-6/STAT3 pathway [58].

In conclusion, it is evident that aging is always accompanied by a decline in probiotics and an increase in harmful bacteria, which is the driving force behind tumorigenesis and cancer progression.

Traditional mechanisms: changes in the immune microenvironment during aging and in tumors

During the process of aging, chronic inflammation, attributed to an imbalance in inflammatory response, is the most distinct variation. Recent studies have revealed that aging-related chronic inflammation can also affect the progression of aging. In *Drosophila*, lower immune deficiency (IMD)/nuclear factor- κ B (NF- κ B) level can mobilize nutrients and prolong lifespan by inhibiting the immune-endocrine axis, whereas higher IMD/NF- κ B levels can result in extensive neurodegeneration and early death by increasing antimicrobial peptides [59]. In humans, C-reactive protein, IL-6, and IL-10 are associated with aging-induced osteoarthritis and cognitive decline in older individuals, presenting as impaired executive function and processing speed [60, 61].

Inflammation not only accelerates the process of aging but also supports cancer cells. In recent decades, researchers have shown the role of inflammation in tumorigenesis and cancer progression. Furthermore, NF- κ B and IL-6 could support the stemness of some cancer cells or induce chemotherapy resistance [62, 63]. NF- κ B-activated cyclooxygenase-2 has been shown to regulate

the magnitude and selectivity of inflammation induced by *Bacillus Calmette–Guérin* in bladder cancer [64]. In addition, tumor necrosis factor- α mediates the interaction between gastric cancer cells and activated fibroblasts, possibly contributing to tumor metastasis [65].

In older individuals, the existence of chronic inflammation also alerts the immune system to immediate danger. During the aging process, every part of the immune system degenerates, which is called immunosenescence. From the perspective of innate immune cells, maturation of natural killer (NK) cells and accurate anti-inflammatory effects of macrophages and neutrophils are apparently impaired in older individuals [66–68]. Considering adaptive immunity, bone marrow and thymic involution lead to reduction in the generation of naïve T/B cells and accumulation of effector memory T/B cells. Moreover, the normal capability of accumulated memory B cells is also damaged [69, 70]. Previous studies have also revealed that age-related chronic persistent inflammation can transform protective immune cells into various immunosuppressive cells, such as regulatory phenotypes of B cells (Bregs), T cells (Tregs), macrophages, and dendritic cells [71].

Many researchers believe that immunosenescence may also be responsible for the genesis and development of tumors. The escape of NK cells in older individuals caused by impaired maturation accelerates the progression of acute myeloid leukemia and decreases the rate of survival [72, 73]. In a separate study on aged mice, subtype remodeling of macrophages, which presented as expanded M2 macrophages in the bone marrow and spleens, was probably carcinogenic [74]. In addition, decreased naïve T cells and increased effector memory T cells would lead to incompetent T cell responses to novel tumor-associated antigens. A previous study on non-small cell lung cancer (NSCLC) showed that CD4+ Tregs were responsible for accelerated tumor growth [75]. Núñez et al. pointed out that cancer invasion in draining lymph nodes was also caused by Treg accumulation [76]. Previous studies have also concluded that tumor immune escape is the consequence of cross-regulation between Bregs and cancer cells [77].

Therefore, the aged systemic microenvironment, especially the accumulation of inflammatory factors and immunosuppressive cells, can accelerate aging anteriorly. This vicious cycle aggravates the imbalance of the inflammatory response and triggers immune escape, which may trigger tumorigenesis.

Energy metabolism and substance metabolism disorders influence both aging and cancer

Metabolism is one of the most basic characteristics of life, and aging is characterized by receding bioenergetics,

which are disorders such as those of lipid synthesis and decomposition and glucose utilization [78]. In the present study, we have summarized the relevance of aging and cancer from the perspectives of systemic energy metabolism and local substance metabolism.

Nicotinamide adenine dinucleotide (NAD⁺) is a key coenzyme that mediates multiple redox reactions and plays an indispensable and irreplaceable role in energy metabolism [79]. Starting from post-puberty, NAD⁺ in both sexes is negatively correlated with age because of increased degradation and decreased generation [80, 81]. In addition to humans, mice and *C. elegans* also show an age-dependent decrease in NAD⁺ [82]. NAD⁺ levels play a crucial role in mitochondrial homeostasis in various species, and aging-mediated absence of NAD⁺ impairs mitochondrial function [83]. Several aging models have shown mitochondrial fission and fusion which can regulate metabolic efficiency to adapt to rapid changes in nutrient availability were impaired in older individuals. Furthermore, decreased mitochondrial trafficking has also been observed in the process of aging [84].

Subsequently, decreased levels of NAD⁺ and mitochondrial dysfunction expedite the aging process. Werner syndrome is a premature aging disease characterized by damaged mitochondria and NAD⁺ deficiency. NAD⁺ repletion can neutralize premature aging symptoms by restoring mitochondrial function and mitophagy [85]. Moreover, failure in NAD⁺ signaling has been proven to have a causal relationship with the vascular [86], immune system [87], and nervous system [88], retina [89], cardiac and skeletal muscles during aging [90]. NAD⁺ can regulate circadian reprogramming to delay aging by suppressing the clock repressor PER2 [91]. However, although NAD⁺ repletion seemed to counteract the aging process, this method could only improve healthspan in normal individuals. Some studies in mice revealed that their lifespan could not be extended or could only be slightly extended (approximately 40 days) by replenishing NAD⁺ levels [92, 93].

In addition, numerous studies have suggested that NAD⁺ and mitochondrial dysfunction are closely related to cancer. The key enzymes of NAD⁺, such as *de novo* synthesis-nicotinate phosphoribosyltransferase and salvage synthesis-nicotinamide phosphoribosyltransferase, are overexpressed in various cancers and contribute to higher glycolytic activity, cancer progression, chemoresistance, and poor prognosis [94–97]. According to Chowdhry et al., the survival of tumors relies on two key enzymes [98]. Moreover, a review summarized the role of NAD⁺ in cancers and concluded that NAD⁺ was the initiator of the Warburg effect in cancers [99]. Therefore, NAD⁺ mainly acts as a carcinogenic factor. Although NAD⁺ is upregulated in cancers, mitochondrial damage and dysfunction such as

increased mitochondrial fission, reduced mitochondrial fusion [100, 101], mutation and depletion of mitochondrial DNA [102-104], disruption of mitochondrial protein [105, 106] and other processes were also universally observed in various cancers which causing cancer growth, metastasis, invasion, and therapeutic resistance.

Regarding local substance metabolism, a previous study conducted by Gomes et al. revealed that the levels of three metabolites—phosphoenolpyruvate (PEP), quinolinate, and methylmalonic acid (MMA) were significantly increased in older individuals compared to those in younger individuals [107]. MMA is a by-product of propionate metabolism that promotes a pro-aggressive epithelial-to-mesenchymal transition-like phenotype by upregulating SOX4 and triggering cancer progression [108]. This may be the most recent cogent evidence to prove the relationship between aging and cancer in terms of substance metabolism. Quinolinate, a metabolite of the kynurenine pathway, participates in NAD⁺ de novo synthesis. Quinolinate phosphoribosyltransferase is the rate-limiting enzyme in the kynurenine pathway, which is overexpressed in breast cancer and promotes cell migration and invasion [109, 110]. In contrast, PEP seems to have anticancer roles. In melanoma, increased PEP bolstered effector functions of tumor-specific T cells, leading to inhibited tumor growth and prolonged survival in rats [111]. This demonstrates that accumulated metabolites during aging may have positive or negative impacts on cancer cells and that the final effect is differ based on tumor types.

In addition, α -ketoglutarate (α -KG), a tricarboxylic acid cycle intermediate, also changes on aging, thus intervening in the processes of both aging and cancer. During the process of aging, Chin et al. reported that α -KG delayed aging and prolonged longevity by approximately 50% by suppressing ATP synthase and decreasing oxygen consumption [112]. The function of α -KG in extending the lifespan has also been observed in *Drosophila* [113]. Last year, Shahmirzadi further proved that replenishing α -KG not only reduced frailty but also prolonged the lifespan of both female and male mice (especially female mice) by inhibiting aging-related chronic inflammation [114]. α -KG can also delay aging-related osteoporosis and fecundity reduction [115, 116].

The effects of α -KG on cancers have also been reported in recent studies. For instance, α -KG supplementation or accumulation can switch glucose metabolism to oxidative phosphorylation from glycolysis, resulting in the inhibition of the Warburg effect, improved tumor suppressors, decreased HIF-1 α levels, or downregulated matrix metalloproteinase 3 to suppress breast cancer oncogenesis, progression, and metastasis [117, 118]. In CRC, dietary α -KG can regulate the

immune system or intestinal flora to inhibit inflammation-related CRC [119]. In contrast, α -KG enhances glucose uptake, sustains cancer cell survival, and accelerates gliomagenesis by activating the IKK β and NF- κ B signaling pathways [120].

Endocrine hormones may delay aging but promote cancer

Aging is accompanied by a recession of the endocrine axis, which contributes to neuroendocrine disequilibrium, manifesting as hormonal absence or excess. Previous studies have reported a reduction in the levels of thyroid stimulating hormone (TSH), growth hormone (GH), luteinizing hormone (LH), and follicle-stimulating hormone (FSH) in older individuals compared to those in younger individuals with the same health status [121-123].

Recently, hormones (TSH, GH, LH, and FSH) have been found to intervene in the lifespan of an individual. Researchers have revealed that the macrobian population is characterized by high TSH generation and TSH insensitivity without altered energy metabolism [124, 125]. Moreover, TSH can reverse aging mediated by BRAFV600E mutations in mice, both in vitro and in vivo [126]. Similar to GH, insulin/insulin-like growth factor I signaling suppression delayed stress granule protein accumulation and subsequently prolonged longevity in *C. elegans* [127, 128]. Furthermore, GH treatment reduced longevity in aged mice (common mice or macrobian mice—Ames dwarfs) by intervening with the genotype, thus confirming the influence of hormones on lifespan [129, 130]. In humans, GH receptor exon 3 deletion homozygotes can prolong the lifespan of males by approximately 10 years [131]. In addition, gonadotropin-releasing hormone, which regulates the FSH/LH ratio, has also been proven to restore impaired neurogenesis and delay aging in mice [132].

The relationship between hormones (TSH, GH, LH, and FSH) and cancer has also been previously reported. Abnormally activated TSH receptors can result in accelerated angiogenesis and deterioration of thyroid cancer [133]. Furthermore, high levels of TSH are associated with an increased risk of thyroid cancer [134]. In addition, TSH can promote the growth of ovarian and liver cancers, resulting in poor survival rates and resistance to chemotherapy [135]. Moreover, GH treatment or receptor stimulation increases cancer incidence and the risk of cancer mortality, induces chemoresistance and metastasis in some specific populations and tumors [136-143]. Notably, people or rodents with developmental GH/IGF-1 show rare-to-no cancer occurrence due to the overexpression of DNA repair genes and perpetual strengthening of DNA repair

capacity [144]. This phenomenon further confirms the powerful effects of GH/IGF-1 on the promotion of cancers. Regarding gonadotropic hormones, FSH upregulates ACTL6A, a novel oncogene that supports glucose metabolism and, thus, increases the invasiveness of ovarian cancer cells [145].

Therefore, aging is always related to endocrine disorders, which can result in an array of symptoms and diseases; hormone replacement therapy occasionally delays aging. However, direct replenishment of hormones is not appropriate because the method appears to be carcinogenic. In conclusion, the present study clearly indicates that hormones are a link between aging and cancer.

Similarity in epigenetic changes between aging and cancers

Epigenetics is a branch of genetics that deals with heritable changes in gene expression without altering the nucleotide sequence. Epigenetic phenomena include DNA methylation, genomic imprinting, maternal effects, gene silencing, and RNA editing. Herein, we have discussed the epigenetic phenomena involved in aging and cancer.

Aging is accompanied by constantly decreasing rates of methylation on a genomic level, which incurs chromosomal instability. Moreover, due to this ascending rate of methylation in CpG island, which results from hypermethylation around the promoter of vital genes, these vital genes will gradually be silenced [146-148]. Recent studies have revealed a close association between DNA methylation age and chronological age. For instance, in embryonic stem cells, the age of DNA methylation was approximately zero, and in centenarians, it was consistent with the venerable age ($r = 0.89$) [149, 150]. Acceleration of the DNA methylation age, in other words, DNA methylation age is older than the chronological age, can improve mortality without being influenced by other risk factors [151]. Furthermore, several studies have revealed that individuals live longer, and their offspring exhibit a deceleration in DNA methylation age [152, 153].

Similarly, acceleration of the DNA methylation age also exists in cancer. Previous studies have shown that epigenetic age in cancer tissues increases by 40% in the same person compared that in normal tissues and that the acceleration was significantly conspicuous by nearly 36.2 years on average [154, 155]. Several studies have proposed that the acceleration of DNA methylation age increases the risk of malignant tumor [156-159]. Furthermore, recent studies have proposed that the level of DNA methylation promoter in body fluids can serve as

an early monitoring index in bladder cancer [160], and lung cancer [161] and some other cancers [162, 163].

Genomic imprinting is characterized by monoallelic expression. A previous study reported that the area of methylation on IGF2 promoters expanded during the aging process and that the initially unmethylated allele was methylated. This study also pointed that similar expansive methylation of IGF2 promoters was also revealed in some cancers, such as colon and lung cancers [164]. In a related study by Fu et al., IGF2 exhibited a tissue-specific loss of imprinting with aging in the prostate lobes of humans and rats, which may also exist in prostate cancer [165]. RasGrf1 is a paternal imprinted gene, and erasure of its imprinting leads to the downregulation of RasGrf1 and prolonged lifespan in mice [166]. Alterations in imprinting of the RasGrf1 gene is also observed in cancer cells. Furthermore, increased methylation of RasGrf1 in the gastric mucosa of patients with gastric cancer has been observed as compared with healthy individuals, which results in a low expression of RasGrf1 as well as tumor invasion [167].

Heterochromatinization is a method of transcriptional gene silencing. In yeast and *Drosophila melanogaster*, loss of gene silencing occurs with age, and this phenomenon results from the loss of repressive heterochromatin during aging [168]. Instability or defects within heterochromatin are widespread alterations in cancer, which also take responsibility for the imbalanced epigenome of tumors [169].

The deepest connection: genetics of aging and cancers

Klotho, a human gene located on chromosome 13, has been dubbed the “master of mortal longevity.” *Klotho* produces two proteins by selective splicing: a membrane protein and a circulating protein. Numerous investigations have established that *Klotho* is closely related to aging. The secretion of *Klotho* has been shown to decrease considerably in the kidneys and serum of aged mice [170]. Additionally, a previous study discovered that between the ages of 21 and 39 years, *Klotho* expression decreases at a rate of 0.082 K/S per year in granulosa cells and 31.95 pg/mL per year in the serum [171].

Recent research has demonstrated that aging-related *Klotho* depletion accelerates aging. Chen discovered that mutations in the *Klotho* gene resulted in the suppression of the transcription factor Nrf2 and inactivation of glutathione reductase, resulting in cardiac aging [170]. Xiaofei Xu et al. demonstrated an intimate relationship between low *Klotho* expression and impaired ovarian reserve [171]. Ullah et al. proposed that *Klotho* deficiency impaired telomerase activity, resulting in stem cell aging and apoptosis [172]. Additionally, it has been demonstrated that the α -*klotho* protein fragment

successfully improves cognitive competence and neural resilience in aged mice [173].

Additionally, *Klotho* acts as a tumor suppressor, disrupting neoplastic glycolysis and mitochondrial function while sparing normal cells [174]. Rubinstein et al. pointed out that treatment with *klotho* or its active region KL1 could inhibit CRC growth in vivo or in vitro by improving the unfolded protein response [175]. *Klotho* also inhibits the NF- κ B pathway and suppresses CCL2 transcription, thereby decreasing CRC progression [176]. *Klotho* functions as an IGF-1R inhibitor in diffuse large B-cell lymphoma, inhibiting cell apoptosis and proliferation [177].

Additionally, there are numerous other common genetic factors associated with aging and cancer. However, it is important to note that some antiaging genes are bidirectional.

AUF1 encodes RNA-binding proteins. Pont et al. postulated that *AUF1* activates the telomerase promoter and assumes responsibility for the maintenance of telomere length. *AUF1*-deficient mice would present with premature aging [178]. Additionally, He et al. confirmed that *AUF1* can prevent aging in vascular endothelial cells [179].

AUF1 has also been shown to affect cancer cells; this regulatory effect is bidirectional. *AUF1* induces apoptosis and suppresses cell proliferation by inhibiting the proto-oncogene *bcl-2* and cyclin D1. Its presence plays a critical role in the antitumor activity of certain tumor suppressors, such as *p63* [180]. In contrast, the overexpression of *AUF1* has been shown to promote malignant transformation and cancer progression [181]. For example, *AUF1* accelerates the development of CRC by activating the ERK1/2 and AKT pathways and is associated with a poor prognosis [182]. Additionally, it promotes the proliferation and invasion of breast cancer and thyroid cancers and results in poor outcomes in patients with hepatocellular carcinoma [183, 184].

The turning point of this bidirectional effect may be the level of expression. In other words, a normal expression of antiaging genes ensures that anticancer pathways remain functional. However, aberrant or obligatory overexpression results in retroaction by excessive activation of tumor promoters.

SIRT1 is downregulated in aging individuals and plays a role in delaying the onset of organic aging [185, 186]. Simultaneously, *SIRT1* acts as a protective gene in certain types of cancers, including gastric and breast cancer, and its deficiency accelerates cancer progression and chemoresistance. Therefore, *SIRT1* overexpression may inhibit the proliferation of cancer cells [187, 188]. However, this protective effect is not universal. In certain cancers, such as CRC and cervical cancer, artificial overexpression significantly enhances proliferation,

invasion, and stemness, whereas spontaneous knock-out has an inhibitory effect [189-191].

Therefore, we may also deduce that the effect of particular antiaging genes on various cancers is highly variable, regardless of whether they are expressed passively or silently, and may differ according to tissue specificity.

p16 is a biochemical indicator of biological senescence that increases with age [192]. Previous studies have established that the downregulation of *p16* expression improves the regenerative capacity of pancreatic β -cells [193] and rescues the degradation of the thymus [194], adipose tissue, skeletal muscle, and eyes [195, 196]. In general, *p16* is a pro-aging element.

In contrast to what we have previously stated, this pro-aging factor generally functions as a tumor suppressor gene. *p16*, a cyclin-dependent kinase inhibitor, arrests cells in the G1 phase. Numerous carcinogens, including miR-30 and constitutive photomorphogenesis 9 signalosome 6, cause cancer by suppressing *p16* expression, enhancing cell division, and interfering with efficient cellular senescence [197, 198]. Liu et al. also demonstrated that ablation of *p16* in B cells could reverse aging while significantly increasing the incidence of B-cell neoplasms [199]. These findings suggest that cellular senescence may have evolved as a strategy to protect against cancer.

In conclusion, from a genetic standpoint, delaying aging does not always suppress cancer.

Aging and cancer have common means of interventions

Antiaging and anticancer agents are two important subjects that compel us to pursue them indefinitely. Here, we summarize the approaches that can delay aging, or even prolong lifespan, and suppress cancer synergistically (Fig. 3, Table 3, and Table 4).

Repurposing common medicines to treat aging and cancer

While researching antiaging and anticancer treatments, researchers surprisingly discovered that some medications possessed dual effects.

1. Metformin.

In mice, metformin delayed ovarian and gut aging, attenuated hearing loss and age-related neurodegeneration, and improved skin healing abilities in aged rats [200-204]. In inbred 129/Sv mice, Anisimov et al. discovered that treatment with metformin reduced the mean lifespan by 13.4% in males, but increased it by 4.4%

in females [205]. However, another study found that the opposite phenomenon of metformin supplementation in the neonatal period extended the mean lifespan (+20%) and maximum life span (+3.5%) in male mice rather than in female mice [206]. Zhu et al. further showed that metformin impaired cardiac homeostasis and longevity in female mice [207]. Therefore, the sex-related differences in the antiaging effects of metformin are significant and warrant further investigation. In humans, metformin delayed the aging of mesenchymal stem cells and reversed mitochondrial function in fibroblasts derived from patients with parkinsonism [208, 209]. Additionally, metformin improves muscle strength and alleviates frailty syndrome in the elderly [210]. In terms of cognitive competence, metformin has been shown to delay

cognitive deterioration in diabetic older adults [211]. An ongoing study sponsored by the Albert Einstein College of Medicine is delving deeper into the antiaging properties of this incredible drug. They compared the number of upregulated genes in the muscle and adipose tissue of young healthy subjects categorized into two groups (metformin treated and placebo treated) after the intervention, which lasted 6 weeks, and discovered that the metformin group had significantly fewer genes than the control group, regardless of muscle or adipose tissue. This finding suggests that metformin may alter biology at the genetic level. By examining the disparate genetic functions, researchers will have a better understanding of the antiaging mechanisms of metformin (<https://clinicaltrials.gov/ct2/show/NCT02432287>).




Figure 3. Common means of interventions between aging and cancer.

To date, numerous studies have been conducted on the anticancer properties of metformin. Metformin has been shown to suppress tumorigenesis, growth, metastasis, and improve chemosensitivity in various cancers in both in vitro and in vivo animal experiments [212-216]. In humans, a recent trial demonstrated that treatment with metformin could improve disease-free survival (not with metformin: HR = 1.40, $p = 0.043$), distant disease-free survival (not with metformin: HR = 1.56, $P = 0.013$), and overall survival (not with metformin: HR = 1.87, $p = 0.004$) in patients with diabetes and HER2-positive breast cancer [217]. Additionally, another phase 2 trial indicated that metformin supplementation increased cisplatin sensitivity and improved overall survival compared to the expected survival in nondiabetic patients with ovarian cancer [218]. Metformin possesses anticancer properties in NSCLC [219]. However, a trial of

patients with advanced pancreatic cancer discovered that the administration of metformin did not improve the prognosis of patients treated with chemotherapy [220]. Therefore, we can deduce that the anti-cancer effects of metformin are unaffected by diabetes mellitus but vary according to tumor type.

Most importantly, Anisimov et al. observed that treatment with metformin initiated at a young and middle-age prolonged longevity (young: +14.1%; middle: +6.1%, $p > 0.05$) and delayed tumorigenesis (young: delay by 22%; middle: delay by 25%) in female SHR mice. Correspondingly, commencing metformin treatment at an advanced age did not increase life expectancy and lost its anticancer effects [221]. Moreover, in 129/Sv mice, metformin extended the mean lifespan of healthy males (+20%, $p < 0.05$) and tumor-bearing males (+4%, although $p = 0.177$). In contrast, it was unable to slow aging in

healthy females, while simultaneously improving survival in tumor-bearing females [206]. These phenomena could be explained in part by the putative associations between

anticancer effects and the antiaging properties of metformin.

Table 3. The effect of different therapies in rats.

Intervention	Prolong lifespan	Inhibit cancer	Both in a model
Medicine			
Metformin	Yes (sex specificity) [205-207]	Yes [212, 215, 216]	Yes [205, 206, 221]
Resveratrol	Yes [226-228]	Yes [236-239]	No evidence
Rapamycin	Yes (sex specificity) [244-246, 351]	Yes [248, 250]	Yes [255, 256, 351-354]
DR			
CR	Yes [264, 265]	Yes [268-270]	Yes [272, 273]
KD	Yes [277]	Yes [279-282]	Yes [277]
Microbiota			
Symbiotic	Only delay aging[286-289]	Yes [295-298]	No evidence
FMT	Yes [302]	Yes [304-307]	No evidence
Metabolites			
NMN	Only delay aging[313-319]	No evidence	No evidence
Spermidine	Yes [327, 330, 331]	Yes [341, 342]	Yes [331]

DR, dietary restriction; CR, caloric restriction; KD, ketogenic diet; FMT, fecal microbiota transplantation; NMN, nicotinamide mononucleotide.

2. Resveratrol

In mice, resveratrol therapy delays aging of the cardiovascular system, improves cognitive performance, and reduces aging-mediated renal impairment [222-225]. Additionally, it increased the life expectancy of HtrA2 knockout mice (mitochondrial dysfunction), SOD1G93A transgenic mice (amyotrophic lateral sclerosis), and SAMP8/1 mice (Alzheimer's disease) [226-228]. In humans, resveratrol treatment has been shown to slow the process of skin aging [229], improve cerebrovascular function, delay cognitive decline [230, 231], and improve mobility-related physical function [232].

The anticancer properties of resveratrol have also been investigated. In an in vitro experiment, resveratrol successfully inhibited the migration, invasion, and viability of cancer cells and induced cell death [233-235]. In vivo experiments showed that resveratrol suppressed tumorigenesis, decreased micro-vessel formation of tumors, alleviated tumor burden, and improved chemotherapeutic response in mice [236-239]. Unfortunately, clear and direct evidence regarding the effects of clinical treatment continues to be insufficient.

3. Rapamycin

Short-term rapamycin treatment inhibited premature ovarian failure, rejuvenated oral health, and slowed the epigenetic age of the liver in aging mouse models [240-242]. In terms of lifespan, Bitto et al. observed that 3 months of rapamycin supplementation increased the life expectancy of middle-aged mice by 60% [243]. Strong et al. further pointed out the sex-specific effects of rapamycin, stating that a 3-month course of rapamycin

improved longevity in male mice, whereas female mice required continuous exposure [244]. However, it has been demonstrated that rapamycin does not extend longevity in progeroid models with DNA repair deficiency or telomerase deletion, regardless of dosage, gender, or timing [245, 246]. This indicates that the antiaging effects of rapamycin are sex-specific and that its mechanism of action may involve enhancing DNA repair capacity and preventing DNA damage. Topical rapamycin treatment delays skin aging and improves skin appearance in humans [247].

In terms of anticancer activity, rapamycin alone or in combination with other agents has been shown to decrease the progression of various tumors or to enhance the chemotherapeutic response in vitro or in vivo [248-250]. Clinically, as an adjuvant therapy, rapamycin successfully developed a pathological response and inhibited metabolic activity in rectal cancer [251], suppressed head and neck tumor growth [252], enhanced the curative effect of intravesical BCG therapy [253], decreased the incidence of secondary skin cancers, and improved squamous cell cancer-free survival at 5 years ($p = 0.007$) in kidney transplant recipients [254]. Of course, these are phase I and II trials, and additional clinical trials are needed to investigate the direct therapeutic effects of rapamycin in various types of cancers.

Several studies on rats have demonstrated a more direct connection between aging and cancer. For example, Anisimov et al. discovered that rapamycin treatment increased the mean (+4.1%) and maximal lifespans (+12.4%), as well as delayed the incidence of breast cancer ($p < 0.001$) and decreased tumor burden ($p < 0.01$) [255]. Wilkinson et al. reported that rapamycin delayed the aging of multiple systems while decreasing the

incidence rates for adrenal tumors ($p = 0.04$) and lung tumors (no statistical significance) [256]. Additionally, researchers discovered that rapamycin, despite its ability to prevent cancer growth, failed to prolong survival in rats already suffering from tumors. Therefore, we conclude

that rapamycin suppresses tumorigenesis by delaying aging rather than delaying death by inhibiting cancer. Hence, there are several targets controlled by rapamycin in cancer and aging.

Table 4. The effect of different therapies in humans.

Intervention	Delay aging	Prolong lifespan ¹	Inhibit cancer	Improve prognosis
Medicine				
Metformin	Yes [208-211]	No evidence	Yes (vitro and vivo) [212-214, 216]	Yes [217-219]
Resveratrol	Yes [229-232]	No evidence	Yes (vitro) [233-235]	No evidence
Rapamycin	Yes [247]	No evidence	Yes (vitro and vivo) [248, 249, 251-254]	No evidence
DR				
CR	Yes [266]	Yes ² [267]	Yes (vivo) ² [271]	No evidence
KD	Yes [278]	No evidence	Yes (vivo) [284]	Yes [284, 285, 355]
Microbiota				
Symbiotic	Yes [290-294]	No evidence	Yes (vivo) ² [299, 301]	Yes ³ [300]
FMT	No evidence	No evidence	Yes (vivo) [308]	Yes [308]
Metabolites				
NMN	No evidence	No evidence	Unclear [320, 321]	No evidence
Spermidine	Yes [332-334]	Yes ² [335]	Yes (vitro and vivo ²) [336-340]	No evidence

DR, dietary restriction; CR, caloric restriction; KD, ketogenic diet; FMT, fecal microbiota transplantation; NMN, nicotinamide mononucleotide.

1: Prolong lifespan means that this method can extend longevity of whole populations rather than patients with specific diseases.

2: Only proved in retrospective study.

3: $0.05 < P\text{-value} < 0.1$

Conserved role of dietary restriction in aging and cancers

In recent years, we have begun to pay more attention to dietary restriction which has been shown to extend the healthy lifespan of several species without causing malnutrition. Caloric restriction (CR), restriction of specific dietary ingredients, ketogenic diet (KD), fasting, and so on are all examples of dietary restriction [257]. In this study, we were mostly concerned about the role of CR and KD in aging and cancer.

1. CR

In rats, CR restored redox homeostasis in aged hearts [258], delayed skeletal muscle degeneration [259], postponed aging-correlative recession in locomotor activity [260], alleviated aging-related metabolic disorders [261], remodeled age-related methylation pattern [262], and even caused epigenetic age deceleration [263]. In terms of longevity, in the last century, people have observed that 40% of long-term CR maintains cellular replication capacity and increases both average lifespan and maximum lifespan by 36% and 20%, respectively [264]. Yan et al. demonstrated that lifelong CR increased the survival of rats by 30% when combined with GH/IGF-1 suppression, while GH/IGF-1 suppression alone only improved survival by 8% [265]. In humans, CR prolonged the replicative lifespan of adipose-derived

stromal/progenitor cells and delayed white adipose tissue recession [266]. A retrospective study revealed that Okinawans who adhered to a CR diet extended both average and maximum longevity compared to Japanese and Americans [267].

In terms of anticancer effects, CR has been shown to decrease the progression of numerous cancers and improve the therapeutic response in mice [268-270]. Additionally, clinical studies have shown that CR successfully reduces the incidence of breast cancer (all women: OR = 0.52, $p = 0.001$; premenopausal women: OR = 0.36, $p < 0.001$; postmenopausal women: OR = 0.77, $p = 0.001$) [271].

The antiaging and anticancer effects of CR were observed using the same models. Weindruch et al. discovered that CR not only extended the lifespan of mice but also suppressed the occurrence of spontaneous lymphoma [272]. Lee et al. found that CR prolonged the maximum lifespan of mice by 13% ($p < 0.0001$) and appeared to decrease the incidence of hepatocellular carcinoma and lymphoma in mice (both $p \leq 0.001$) [273].

2. KD

KD is a low-carbohydrate diet with a high fat content (5–10% of the total daily consumed kcal). Recently, it was discovered that non-obesogenic KD improved memory [274], inhibited myocardial remodeling and damage [275], and preserved skeletal muscle mass and function in

mice during the aging process [276]. Additionally, Megan et al. demonstrated that KD preserved physiological function while increasing median longevity by 13.6% along with survival in mice [277]. Numerous investigations conducted in humans have established that KD improves cognitive function and memory in patients with Alzheimer's disease, particularly those with the APOE4 genotype, even in short-term memory [278].

The anticancer effects of KD were also investigated. For example, in a mouse model, KD alone has been shown to inhibit the growth of a variety of tumors [279, 280]. In mice with pancreatic cancer, combining radiotherapy with KD may improve radiation sensitivity and extend survival time compared to radiotherapy alone [281]. Additionally, it may enhance the anticancer properties of rapamycin and other chemotherapeutic agents [279, 282]. In humans, KD has been shown to enhance the effects of chemotherapy, promote the incidence of complete or partial responses, and increase patient survival rate without inducing side effects [283-285]. In summary, KD serves as an adjuvant therapy to enhance the effectiveness of conventional therapy.

It is worth mentioning that KD not only increased median lifespan by 13.6% in a mouse model, but also decreased the incidence of malignancies, particularly histiocytic sarcoma ($P < 0.1$), at the time of death. This effect was dose-dependent [277]. This result further established that energy and substance metabolism connect aging with cancer, which is likely to be mediated by mitochondrial function and oxidative stress.

Emerging approaches: interventions in the gut microbiota in aging and cancer

Previous research has established that disturbances in the intestinal microbiota occur inexorably with aging and can result in various age-related diseases, including cancer. Thus, microbiota intervention strategies such as probiotics, prebiotics, symbiotics, and fecal microbiota transplantation (FMT) are emerging to rectify flora disturbances.

1. Symbiotic

Symbiotic bacteria have been shown to increase cognitive function, minimize memory deterioration, delay behavioral degeneration (such as muscular vigor and exploratory activity), and protect against primary and secondary osteoporosis in aging mice [286-289]. In humans, earlier research concluded that symbiotic bacteria restore the skin barrier; protect against photoaging; and improve hair quality [290], cognitive function, and aging-mediated memory impairment during aging [291-293]. Liu et al. also argued that ingesting

symbiotics significantly upregulated serum calcium concentrations in older individuals, although its benefits to bone health remained ambiguous [294].

Recently, a plethora of evidence supporting the anticancer effects of symbiotics has emerged. Previous research has shown that symbiotics, such as *Bifidobacterium*, *Bacteroides fragilis*, and *Akkermansia muciniphila*, can inhibit cancer progression, mainly breast and CRC, and enhance chemotherapy effects in vitro or in mice [295-298]. Clinically, Toi et al. discovered that *Lactobacillus casei strain Shirota* consumption showed a negative correlation with breast cancer incidence (OR = 0.65, $p = 0.048$) [299]. Takada et al. demonstrated that supplementation with probiotics improved progression-free survival (HR = 1.73, $p = 0.0229$), disease control (OR = 0.51, $p = 0.0004$), and overall response (OR = 0.43, $p < 0.0001$) in NSCLC patients receiving anti-programmed cell death 1 protein (anti-PD-1) monotherapy [300]. Additionally, the use of probiotics decreased the incidence of metachronous gastric cancer after endoscopic resection (HR = 0.29, $p = 0.034$) [301].

2. FMT

It is worth noting that FMT from wild type mice prolongs the life of progeroid mice, and it has been established that *Akkermansia muciniphila* plays a predominant role. Antiaging effects may be dependent on secondary bile acid restoration [302]. Additionally, another study discovered that FMT was capable of successfully regulating Hippo signaling, which is involved in the manipulation of aging processes [303].

An earlier study showed that FMT from normal mice to tumor-bearing mice decreased tumor load and size in the azoxymethane-dextran sodium sulfate (AOM-DSS) model of CRC [304, 305]. Gopalakrishnan et al. observed that the gut microbiome of patients responding to anti-PD-1 could improve antitumor immunity in germ-free mice and restore the effects of PD-1 blockade in resistant mice [306, 307]. In humans, a recent clinical study discovered that FMT from healthy obese individuals to patients with gastroesophageal cancer improved therapeutic response ($p = 0.035$), and may also improve overall survival (HR = 0.38, $p = 0.057$) and progression-free survival (HR = 0.5, $p = 0.092$) in patients [308].

The use of gut microbiota intervention for anti-aging and anti-cancer strategies is currently limited. Additional evidence is necessary to support this therapy.

Replenishing insufficient metabolites: is it effective and safe enough?

Previous research has shown that circulatory metabolites, including NAD⁺, glutathione, spermidine, glutamine, and

α -ketoglutarate, decrease significantly in the older individuals [309-312]. Therefore, the role of restoring these depleted metabolites in antiaging processes is of interest. Here, we have focused on nicotinamide mononucleotide (NMN) and spermidine.

1. NMN

NMN, as an NAD⁺ booster, has been shown to delay vascular aging, promote functional vascular rejuvenation, and reverse vascular dysfunction in aged mice [313, 314]. It also alleviates memory recession and improves learning and cognitive function in mice [315-317]. Additionally, NMN supplementation has been shown to enhance osteogenesis and improve bone density, locomotor activity, and eye function in aged mice [318, 319]. However, the direct antiaging effects of NMN have not been elucidated in humans yet.

With regard to cancer, NAD⁺ is catalyzed by nicotinamide mononucleotide adenylyl transferases (NMNATs, the rate-limiting enzymes) from NMN. In humans, earlier research found that NMNAT2 is highly expressed in CRC and its increased expression correlates with a deeper invasive depth and an advanced TNM stage ($p < 0.05$) [320]. Additionally, as previously stated, NAD⁺ is a known carcinogen in a wide variety of malignancies. NMN supplementation appears to be involved in the progression of cancer. However, Grozio et al. discovered that increased expression of SLC12A8, a particular NMN transporter required for NMN uptake, was associated with a favorable prognosis in patients with pancreatic and breast cancer [321]. Zong et al. discovered that in a mouse model, NMN supplementation could prevent liver fibrosis, which could progress to liver cancer [322].

However, the effects of NMN on cancers are unknown and it is unclear whether replenishing NMN for antiaging purposes is carcinogenic.

2. Spermidine

Recent research in aged mice discovered that dietary spermidine can improve structural brain measures [323], improve spatial and temporal memory [324], decrease locomotor activity loss [325], protect cardiac function [326, 327], delay skeletal muscle atrophy [328] and reverse arterial aging [329]. In terms of lifetime, it has been observed that dietary polyamines (spermine and spermidine) prolong longevity and improve survival in ICR mice [330]. Eisenberg et al. further reported that late-life spermidine replenishment (which is easier to achieve in humans) successfully improved the median lifespan by 10% [327]. Yue et al. pointed out that lifelong spermidine supplementation can increase the lifespan of mice by 25%

[331]. In humans, spermidine has been shown to preserve female fertility [332], restore skin structure and barrier function [333] and improve mnemonic discrimination performance [334]. In humans, spermidine has been shown to preserve female fertility [335].

Similarly, spermidine was found to decrease cervical cancer in vitro by suppressing HeLa cell proliferation and inducing apoptosis intrinsically or via an apoptosis-inducing factor [336-338]. Razvi et al. discovered that acylspermidine derivatives inhibited breast and blood cancer cell proliferation and triggered apoptosis [339]. Additionally, the presence of spermine/spermidine could induce apoptosis in CRC cells mediated by maize polyamine oxidase [340]. In mice, spermidine inhibited the growth of CT26 CRC cells transplanted into immunocompetent individuals [341]. Soda et al. also discovered that spermine and spermidine intake decreased the incidence of colon cancer induced by a chemical carcinogen in BALB/c mice [342]. Vargas et al. discovered that spermine and spermidine could significantly decrease the risk of CRC in women with a body mass index of ≤ 25 (HR = 0.58, $p = 0.012$) or in women who consume more fiber than the average (HR = 0.44, $p = 0.015$) [343].

Most importantly, Yue et al. demonstrated that spermidine not only extended the lifespan but also alleviated liver fibrosis and inhibited hepatocellular carcinoma by improving MAP1S-mediated autophagy in the same model. This phenomenon indicated that spermidine could regulate both aging and cancer synergistically, possibly by manipulating autophagy [331].

Discussion

Aging is an inevitable process. Economic, technological, and medical advancements have resulted in a decline in birth rates and an increase in the proportion of the older individuals, putting a strain on the health system. The increased incidence of various diseases in aged individuals implies that aging impairs health. As one of the leading causes of death among the elderly, a higher diagnostic rate of malignant tumors in the elderly also suggests a possible link between cancer and aging. For example, in the United States, more than 60% of all lung cancer cases occurred in those aged more than 65 years, whereas less than 2% of lung cancer occurred in people younger than 45 years [344]. Hospitalization costs associated with anticancer treatments have increased significantly in recent years. To extend a healthy lifespan and alleviate the strain on the health care system, it is critical to explore the link between aging and cancer. With the advancement of scientific research, significant changes in the human body during the process of aging

have already been identified, including chronic inflammation, immunosuppression, and endocrine disorders. Additionally, numerous pathogenic mechanisms for common cancers have been proposed, and we identified some aging-related alterations associated with tumor occurrence. Worse yet, these age-related alterations might accelerate the rate of aging, creating a vicious circle. Hence, cancer therapy targeting aging is likely to be effective.

During the last century, researchers have already noticed a decreased incidence of malignant tumors in patients with Huntington's disease, which may be attributed to accelerated programmed cell death of precancerous cells [345]. Several years ago, Musicco et al. discovered that older individuals with Alzheimer's disease had a lower risk of developing malignant tumors [346]. These findings appear to support a negative correlation between cancer, proliferative disease, and aging, which is typically associated with degenerative diseases.

However, this correlation was not statistically significant. Last year, Ording et al. conducted additional research on the correlation between Alzheimer's disease and cancer. They established that the negative correlation between these two diseases was quite limited (SIR = 0.94, 95% confidence interval 0.92–0.96) and weakened over time [347]. Freedman et al. found no correlation between site-specific cancers and amyotrophic lateral sclerosis [348]. Liu et al. discovered a statistically significant positive correlation between the occurrence of melanoma and Parkinson's disease [349].

The association between cancer and degenerative diseases does not always exhibit an inverse correlation; it can occasionally exhibit a positive correlation. This suggests that the relationships between cancer and aging, or organic aging, are intricate, and their tissue specificity should be investigated.

In this review, we summarized numerous strategies for delaying aging and restoring aging-related imbalances or disorders. Additionally, these techniques have been shown to have anticancer effects. At the same time, we further summarized the experimental phases of these strategies to estimate their feasibility and reliability.

Metformin is a relatively well-studied medication. Researchers have established that it can slow organic aging and ameliorate the symptoms of senile diseases. It may also act as a regulator of human genes, bringing their expression levels closer to those of young people (<https://clinicaltrials.gov/ct2/show/NCT02432287>).

Cancer has the potential to suppress tumor progression both in vivo and in vitro, as well as improve patient prognosis. However, its anticancer or antiaging effects may be gender- and tissue-specific, which have not been adequately investigated. Resveratrol and rapamycin

have also been shown to delay organic aging and slow tumor progression; however, evidence of improved prognosis in cancer patients is currently lacking. Dietary restriction has already been shown to have anticancer and antiaging properties, which has been partially verified in humans. Additionally, there is a significant problem in that the level (percentage) or pattern (persistent or intermittent) of restriction, the species of restricted energy, and age or gender disparity are ambiguous. The effects of microbiota intervention on aging and cancer have been primarily tested in animals. Owing to concerns regarding the complexity of the methods, the survival rate of xenogenous microflora, and demic tolerance, the application of FMT in the human body is rather uncommon. Thus, its antiaging and anticancer effects must be further validated. Replenishment of scarce metabolites in elderly individuals, such as spermidine, is another attempt at this monumental task that has been demonstrated in animals to a certain extent. However, clinical trials in humans have not yet begun. Additionally, researchers discovered that spermidine, a polyamine, may suppress carcinogenesis but deteriorates established tumors [350]. Its bidirectional characteristics remain obscure. Additionally, all the approaches outlined above lack direct evidence of extending healthy human life and can only be used as adjuvant medications or therapies for cancer treatment.

Hence, one of the most critical objectives of future research will be to optimize interventions for them to be extensively employed in clinical practice. For example, extending the scope of clinical trials based on safety to further evaluate the therapeutic effects and determine the optimal dose. Developing an appropriate dosage or compound that maximizes bioavailability, minimizes side effects, and improves compliance is critical. Additionally, determining the primary and most appropriate target for each individual and developing an individualized treatment plan are necessary before being extensively used in the clinic. Senolysis offers an alternative to the traditional prevention paradigm of "antiaging therapy," which entails lifestyle or pharmacological interventions to delay the onset of age-induced decrepitude and the occurrence of diseases. The complexity of the human body and the diversity of aging and cancer mechanisms prevent us from conducting extensive and exhaustive research. Thus, additional methods need to be explored. It is notable that blind supplementation with missing substances is not a good idea. In contrast, it may fail to delay aging and even induce tumorigenesis. In the future, we should devote our efforts to precisely identify therapies that would effectively prevent the incidence and progression of cancer by delaying aging.

References

- [1] Hong Y C (2013). Aging society and environmental health challenges. *Environ Health Perspect*, 121(3):A68-A69.
- [2] Song P, Chen Y (2015). Public policy response, aging in place, and big data platforms: Creating an effective collaborative system to cope with aging of the population. *Biosci Trends*, 9(1):1-6.
- [3] Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G (2013). The hallmarks of aging. *Cell*, 153(6):1194-1217.
- [4] Marosi C, Koller M (2016). Challenge of cancer in the elderly. *ESMO Open*, 1(3):e20.
- [5] Deng F, Xu X, Lv M, Ren B, Wang Y, Guo W, et al. (2017). Age is associated with prognosis in serous ovarian carcinoma. *J Ovarian Res*, 10(1):36.
- [6] Migkou M, Kastritis E, Roussou M, Gkotsamanidou M, Gavriatopoulou M, Nikitas N, et al. (2011). Short progression-free survival predicts for poor overall survival in older patients with multiple myeloma treated upfront with novel agent-based therapy. *Eur J Haematol*, 87(4):323-329.
- [7] Gomes AP, Ilter D, Low V, Endress JE, Fernandez-Garcia J, Rosenzweig A, et al. (2020). Age-induced accumulation of methylmalonic acid promotes tumour progression. *Nature*, 585(7824):283-287.
- [8] Calcinotto A, Kohli J, Zagato E, Pellegrini L, Demaria M, Alimonti A (2019). Cellular Senescence: Aging, Cancer, and Injury. *Physiol Rev*, 99(2):1047-1078.
- [9] Preto A, Singhrao SK, Houghton MF, Kipling D, Wynford-Thomas D, Jones CJ (2004). Telomere erosion triggers growth arrest but not cell death in human cancer cells retaining wild-type p53: implications for antitelomerase therapy. *Oncogene*, 23(23):4136-4145.
- [10] Hayes JD, Dinkova-Kostova AT, Tew KD (2020). Oxidative Stress in Cancer. *Cancer Cell*, 38(2):167-197.
- [11] Courtois-Cox S, Jones SL, Cichowski K (2008). Many roads lead to oncogene-induced senescence. *Oncogene*, 27(20):2801-2809.
- [12] Garbers C, Kuck F, Aparicio-Siegmund S, Konzak K, Kessenbrock M, Sommerfeld A, et al. (2013). Cellular senescence or EGFR signaling induces Interleukin 6 (IL-6) receptor expression controlled by mammalian target of rapamycin (mTOR). *Cell Cycle*, 12(21):3421-3432.
- [13] Peeper DS, Shvarts A, Brummelkamp T, Douma S, Koh EY, Daley GQ, et al. (2002). A functional screen identifies hDRIL1 as an oncogene that rescues RAS-induced senescence. *Nat Cell Biol*, 4(2):148-153.
- [14] Brehme M, Voisine C, Rolland T, Wachi S, Soper JH, Zhu Y, et al. (2014). A chaperone subnetwork safeguards proteostasis in aging and neurodegenerative disease. *Cell Rep*, 9(3):1135-1150.
- [15] Kaushik S, Cuervo AM (2015). Proteostasis and aging. *Nat Med*, 21(12):1406-1415.
- [16] Mejias-Pena Y, Rodriguez-Miguel P, Fernandez-Gonzalo R, Martinez-Florez S, Almar M, de Paz JA, et al. (2016). Effects of aerobic training on markers of autophagy in the elderly. *Age (Dordr)*, 38(2):33.
- [17] Phadwal K, Alegre-Abarrategui J, Watson AS, Pike L, Anbalagan S, Hammond EM, et al. (2012). A novel method for autophagy detection in primary cells: impaired levels of macroautophagy in immunosenescent T cells. *Autophagy*, 8(4):677-689.
- [18] Escobar KA, Cole NH, Mermier CM, VanDusseldorp T A (2019). Autophagy and aging: Maintaining the proteome through exercise and caloric restriction. *Aging Cell*, 18(1):e12876.
- [19] Kelmer SE, Kirkpatrick JM, Mazzetto M, Baumgart M, Bartolome A, Di Sanzo S, et al. (2020). Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol*, 16(6):e9596.
- [20] Koziel R, Greussing R, Maier AB, Declercq L, Jansen-Durr P (2011). Functional interplay between mitochondrial and proteasome activity in skin aging. *J Invest Dermatol*, 131(3):594-603.
- [21] Kruegel U, Robison B, Dange T, Kahlert G, Delaney JR, Kotireddy S, et al. (2011). Elevated proteasome capacity extends replicative lifespan in *Saccharomyces cerevisiae*. *Plos Genet*, 7(9):e1002253.
- [22] Tonoki A, Kuranaga E, Tomioka T, Hamazaki J, Murata S, Tanaka K, et al. (2009). Genetic evidence linking age-dependent attenuation of the 26S proteasome with the aging process. *Mol Cell Biol*, 29(4):1095-1106.
- [23] Bader M, Benjamin S, Wapinski OL, Smith DM, Goldberg AL, Steller H (2011). A conserved F box regulatory complex controls proteasome activity in *Drosophila*. *Cell*, 145(3):371-382.
- [24] Moskalev A, Shaposhnikov M, Turysheva E (2009). Life span alteration after irradiation in *Drosophila melanogaster* strains with mutations of Hsf and Hsps. *Biogerontology*, 10(1):3-11.
- [25] Crowe E, Sell C, Thomas JD, Johannes GJ, Torres C (2009). Activation of proteasome by insulin-like growth factor-I may enhance clearance of oxidized proteins in the brain. *Mech Ageing Dev*, 130(11-12):793-800.
- [26] Chondrogianni N, Sakellari M, Lefaki M, Papaevgeniou N, Gonos E S (2014). Proteasome activation delays aging in vitro and in vivo. *Free Radic Biol Med*, 71(303-320).
- [27] Chondrogianni N, Petropoulos I, Franceschi C, Friguet B, Gonos ES (2000). Fibroblast cultures from healthy centenarians have an active proteasome. *Exp Gerontol*, 35(6-7):721-728.
- [28] Hansen M, Rubinsztein DC, Walker DW (2018). Autophagy as a promoter of longevity: insights from model organisms. *Nat Rev Mol Cell Biol*, 19(9):579-593.
- [29] Kroll J (2005). Chaperones and longevity. *Biogerontology*, 6(5):357-361.
- [30] Ito A, Zhao Q, Tanaka Y, Yasui M, Katayama R, Sun S, et al. (2021). Metolazone upregulates mitochondrial chaperones and extends lifespan in

- Caenorhabditis elegans. *Biogerontology*, 22(1):119-131.
- [31] Tong L, Shen S, Huang Q, Fu J, Wang T, Pan L, et al. (2020). Proteasome-dependent degradation of Smad7 is critical for lung cancer metastasis. *Cell Death Differ*, 27(6):1795-1806.
- [32] Guo S, Chen Y, Yang Y, Zhang X, Ma L, Xue X, et al. (2021). TRIB2 modulates proteasome function to reduce ubiquitin stability and protect liver cancer cells against oxidative stress. *Cell Death Dis*, 12(1):42.
- [33] Yu T, Guo F, Yu Y, Sun T, Ma D, Han J, et al. (2017). *Fusobacterium nucleatum* Promotes Chemoresistance to Colorectal Cancer by Modulating Autophagy. *Cell*, 170(3):548-563.
- [34] Yamamoto K, Venida A, Yano J, Biancur DE, Kakiuchi M, Gupta S, et al. (2020). Autophagy promotes immune evasion of pancreatic cancer by degrading MHC-I. *Nature*, 581(7806):100-105.
- [35] Zhang Y, Wang J, Wang X, Gao T, Tian H, Zhou D, et al. (2020). The autophagic-lysosomal and ubiquitin proteasome systems are simultaneously activated in the skeletal muscle of gastric cancer patients with cachexia. *Am J Clin Nutr*, 111(3):570-579.
- [36] Hoter A, Rizk S, Naim HY (2019). The Multiple Roles and Therapeutic Potential of Molecular Chaperones in Prostate Cancer. *Cancers (Basel)*, 11(8):
- [37] Clements SJ (2018). Diet, the intestinal microbiota, and immune health in aging. *Crit Rev Food Sci Nutr*, 58(4):651-661.
- [38] Mitsuoka T (2014). Establishment of intestinal bacteriology. *Biosci Microbiota Food Health*, 33(3):99-116.
- [39] Rubio C, Lizarraga E, Alvarez-Cilleros D, Perez-Pardo P, Sanmartin-Salinas P, Toledo-Lobo M V, et al. (2020). Aging in male Wistar rats associates with changes in intestinal microbiota, gut structure and cholecystokinin-mediated gut-brain axis function. *J Gerontol A Biol Sci Med Sci*,
- [40] Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turroni S, et al. (2016). Gut Microbiota and Extreme Longevity. *Curr Biol*, 26(11):1480-1485.
- [41] Rampelli S, Soverini M, D'Amico F, Barone M, Tavella T, Monti D, et al. (2020). Shotgun Metagenomics of Gut Microbiota in Humans with up to Extreme Longevity and the Increasing Role of Xenobiotic Degradation. *Msystems*, 5(2):
- [42] Barcena C, Valdes-Mas R, Mayoral P, Garabaya C, Durand S, Rodriguez F, et al. (2019). Healthspan and lifespan extension by fecal microbiota transplantation into progeroid mice. *Nat Med*, 25(8):1234-1242.
- [43] van der Lugt B, van Beek AA, Aalvink S, Meijer B, Sovran B, Vermeij W P, et al. (2019). *Akkermansia muciniphila* ameliorates the age-related decline in colonic mucus thickness and attenuates immune activation in accelerated aging *Ercc1 (-/Delta7)* mice. *Immun Ageing*, 16(6).
- [44] Wang T, Cai G, Qiu Y, Fei N, Zhang M, Pang X, et al. (2012). Structural segregation of gut microbiota between colorectal cancer patients and healthy volunteers. *Isme J*, 6(2):320-329.
- [45] Guo S, Chen J, Chen F, Zeng Q, Liu W L, Zhang G (2020). Exosomes derived from *Fusobacterium nucleatum*-infected colorectal cancer cells facilitate tumour metastasis by selectively carrying miR-1246/92b-3p/27a-3p and CXCL16. *Gut*,
- [46] Chen S, Su T, Zhang Y, Lee A, He J, Ge Q, et al. (2020). *Fusobacterium nucleatum* promotes colorectal cancer metastasis by modulating KRT7-AS/KRT7. *Gut Microbes*, 11(3):511-525.
- [47] Parhi L, Alon-Maimon T, Sol A, Nejman D, Shhadeh A, Fainsod-Levi T, et al. (2020). Breast cancer colonization by *Fusobacterium nucleatum* accelerates tumor growth and metastatic progression. *Nat Commun*, 11(1):3259.
- [48] Chen S, Su T, Zhang Y, Lee A, He J, Ge Q, et al. (2020). *Fusobacterium nucleatum* promotes colorectal cancer metastasis by modulating KRT7-AS/KRT7. *Gut Microbes*, 11(3):511-525.
- [49] Hashemi GN, Heidarzadeh S, Jahangiri S, Farhood B, Mortezaee K, Khanlarkhani N, et al. (2019). *Fusobacterium nucleatum* and colorectal cancer: A mechanistic overview. *J Cell Physiol*, 234(3):2337-2344.
- [50] Lopes A, Billard E, Casse AH, Villeger R, Veziat J, Roche G, et al. (2020). Colibactin-positive *Escherichia coli* induce a procarcinogenic immune environment leading to immunotherapy resistance in colorectal cancer. *Int J Cancer*, 146(11):3147-3159.
- [51] Bao Y, Tang J, Qian Y, Sun T, Chen H, Chen Z, et al. (2019). Long noncoding RNA BFAL1 mediates enterotoxigenic *Bacteroides fragilis*-related carcinogenesis in colorectal cancer via the RHEB/mTOR pathway. *Cell Death Dis*, 10(9):675.
- [52] Butt J, Jenab M, Werner J, Fedirko V, Weiderpass E, Dahm CC, et al. (2021). Association of Pre-diagnostic Antibody Responses to *Escherichia coli* and *Bacteroides fragilis* Toxin Proteins with Colorectal Cancer in a European Cohort. *Gut Microbes*, 13(1):1-14.
- [53] Fan L, Qi Y, Qu S, Chen X, Li A, Hendi M, et al. (2021). *B. adolescentis* ameliorates chronic colitis by regulating Treg/Th2 response and gut microbiota remodeling. *Gut Microbes*, 13(1):1-17.
- [54] Parisa A, Roya G, Mahdi R, Shabnam R, Maryam E, Malihe T (2020). Anti-cancer effects of *Bifidobacterium* species in colon cancer cells and a mouse model of carcinogenesis. *Plos One*, 15(5):e232930.
- [55] Heydari Z, Rahaie M, Alizadeh AM, Agah S, Khalighfard S, Bahmani S (2019). Effects of *Lactobacillus acidophilus* and *Bifidobacterium bifidum* Probiotics on the Expression of MicroRNAs 135b, 26b, 18a and 155, and Their Involving Genes in Mice Colon Cancer. *Probiotics Antimicrob Proteins*, 11(4):1155-1162.
- [56] Fahmy CA, Gamal-Eldeen AM, El-Hussieny EA, Raafat BM, Mehanna NS, Talaat RM, et al. (2019). *Bifidobacterium longum* Suppresses Murine Colorectal Cancer through the Modulation of

- oncomiRs and Tumor Suppressor miRNAs. *Nutr Cancer*, 71(4):688-700.
- [57] Agah S, Alizadeh AM, Mosavi M, Ranji P, Khavari-Daneshvar H, Ghasemian F, et al. (2019). More Protection of *Lactobacillus acidophilus* Than *Bifidobacterium bifidum* Probiotics on Azoxymethane-Induced Mouse Colon Cancer. *Probiotics Antimicrob Proteins*, 11(3):857-864.
- [58] Ma J, Sun L, Liu Y, Ren H, Shen Y, Bi F, et al. (2020). Alter between gut bacteria and blood metabolites and the anti-tumor effects of *Faecalibacterium prausnitzii* in breast cancer. *Bmc Microbiol*, 20(1):82.
- [59] Kounatidis I, Chtarbanova S, Cao Y, Hayne M, Jayanth D, Ganetzky B, et al. (2017). NF-kappaB Immunity in the Brain Determines Fly Lifespan in Healthy Aging and Age-Related Neurodegeneration. *Cell Rep*, 19(4):836-848.
- [60] Guan Y J, Li J, Yang X, Du S, Ding J, Gao Y, et al. (2018). Evidence that miR-146a attenuates aging- and trauma-induced osteoarthritis by inhibiting Notch1, IL-6, and IL-1 mediated catabolism. *Aging Cell*, 17(3):e12752.
- [61] Tegeler C, O'Sullivan JL, Bucholtz N, Goldeck D, Pawelec G, Steinhagen-Thiessen E, et al. (2016). The inflammatory markers CRP, IL-6, and IL-10 are associated with cognitive function--data from the Berlin Aging Study II. *Neurobiol Aging*, 38(112-117).
- [62] Sun B, Han Y, Cai H, Huang H, Xuan Y (2021). Long non-coding RNA SNHG3, induced by IL-6/STAT3 transactivation, promotes stem cell-like properties of gastric cancer cells by regulating the miR-3619-5p/ARL2 axis. *Cell Oncol (Dordr)*, 44(1):179-192.
- [63] Wang L, Guo J, Zhou J, Wang D, Kang X, Zhou L (2020). NF-kappaB maintains the stemness of colon cancer cells by downregulating miR-195-5p/497-5p and upregulating MCM2. *J Exp Clin Cancer Res*, 39(1):225.
- [64] Ibrahim OM, Basse P H, Jiang W, Guru K, Chatta G, Kalinski P (2021). NFkappaB-Activated COX2/PGE2/EP4 Axis Controls the Magnitude and Selectivity of BCG-Induced Inflammation in Human Bladder Cancer Tissues. *Cancers (Basel)*, 13(6):
- [65] Zhou Q, Wu X, Wang X, Yu Z, Pan T, Li Z, et al. (2020). The reciprocal interaction between tumor cells and activated fibroblasts mediated by TNF-alpha/IL-33/ST2L signaling promotes gastric cancer metastasis. *Oncogene*, 39(7):1414-1428.
- [66] Jackaman C, Tomay F, Duong L, Abdol RN, Pixley FJ, Metharom P, et al. (2017). Aging and cancer: The role of macrophages and neutrophils. *Ageing Res Rev*, 36(105-116).
- [67] Sapey E, Greenwood H, Walton G, Mann E, Love A, Aaronson N, et al. (2014). Phosphoinositide 3-kinase inhibition restores neutrophil accuracy in the elderly: toward targeted treatments for immunosenescence. *Blood*, 123(2):239-248.
- [68] Lopez-Sejas N, Campos C, Hassouneh F, Sanchez-Correa B, Tarazona R, Pera A, et al. (2016). Effect of CMV and Aging on the Differential Expression of CD300a, CD161, T-bet, and Eomes on NK Cell Subsets. *Front Immunol*, 7(476).
- [69] Bulati M, Caruso C, Colonna-Romano G (2017). From lymphopoiesis to plasma cells differentiation, the age-related modifications of B cell compartment are influenced by "inflamm-aging". *Ageing Res Rev*, 36(125-136).
- [70] Alves A S, Bueno V (2019). Immunosenescence: participation of T lymphocytes and myeloid-derived suppressor cells in aging-related immune response changes. *Einstein (Sao Paulo)*, 17(2):B4733.
- [71] Salminen A (2020). Activation of immunosuppressive network in the aging process. *Ageing Res Rev*, 57(100998).
- [72] Manser AR, Uhrberg M (2016). Age-related changes in natural killer cell repertoires: impact on NK cell function and immune surveillance. *Cancer Immunol Immunother*, 65(4):417-426.
- [73] Sanchez-Correa B, Campos C, Pera A, Bergua JM, Arcos MJ, Banas H, et al. (2016). Natural killer cell immunosenescence in acute myeloid leukaemia patients: new targets for immunotherapeutic strategies? *Cancer Immunol Immunother*, 65(4):453-463.
- [74] Jackaman C, Radley-Crabb H G, Soffe Z, Shavlakadze T, Grounds MD, Nelson DJ (2013). Targeting macrophages rescues age-related immune deficiencies in C57BL/6J geriatric mice. *Aging Cell*, 12(3):345-357.
- [75] Alvisi G, Brummelman J, Puccio S, Mazza EM, Tomada EP, Losurdo A, et al. (2020). IRF4 instructs effector Treg differentiation and immune suppression in human cancer. *J Clin Invest*, 130(6):3137-3150.
- [76] Nunez NG, Tosello BJ, Ramos RN, Richer W, Cagnard N, Anderfuhren CD, et al. (2020). Tumor invasion in draining lymph nodes is associated with Treg accumulation in breast cancer patients. *Nat Commun*, 11(1):3272.
- [77] Shang J, Zha H, Sun Y (2020). Phenotypes, Functions, and Clinical Relevance of Regulatory B Cells in Cancer. *Front Immunol*, 11(582657).
- [78] (2017). Aging and Metabolism: Two Sides of the Same Coin. *Ebiomedicine*, 21:1-2.
- [79] Navas LE, Camero A (2021). NAD(+) metabolism, stemness, the immune response, and cancer. *Signal Transduct Target Ther*, 6(1):2.
- [80] Massudi H, Grant R, Braidy N, Guest J, Farnsworth B, Guillemin GJ (2012). Age-associated changes in oxidative stress and NAD+ metabolism in human tissue. *Plos One*, 7(7):e42357.
- [81] Garrido A, Djouder N (2017). NAD(+) Deficits in Age-Related Diseases and Cancer. *Trends Cancer*, 3(8):593-610.
- [82] Fang E F, Scheibye-Knudsen M, Brace L E, Kassahun H, SenGupta T, Nilsen H, et al. (2014). Defective mitophagy in XPA via PARP-1 hyperactivation and NAD(+)/SIRT1 reduction. *Cell*, 157(4):882-896.
- [83] Katsyuba E, Mottis A, Zietak M, De Franco F, van der Velpen V, Gariani K, et al. (2018). De novo NAD(+)

- synthesis enhances mitochondrial function and improves health. *Nature*, 563(7731):354-359.
- [84] Sharma A, Smith H J, Yao P, Mair WB (2019). Causal roles of mitochondrial dynamics in longevity and healthy aging. *Embo Rep*, 20(12):e48395.
- [85] Fang EF, Hou Y, Lautrup S, Jensen MB, Yang B, SenGupta T, et al. (2019). NAD(+) augmentation restores mitophagy and limits accelerated aging in Werner syndrome. *Nat Commun*, 10(1):5284.
- [86] Das A, Huang GX, Bonkowski MS, Longchamp A, Li C, Schultz MB, et al. (2019). Impairment of an Endothelial NAD(+)-H2S Signaling Network Is a Reversible Cause of Vascular Aging. *Cell*, 176(4):944-945.
- [87] Minhas PS, Liu L, Moon PK, Joshi AU, Dove C, Mhatre S, et al. (2019). Macrophage de novo NAD(+) synthesis specifies immune function in aging and inflammation. *Nat Immunol*, 20(1):50-63.
- [88] Lautrup S, Sinclair DA, Mattson MP, Fang EF (2019). NAD(+) in Brain Aging and Neurodegenerative Disorders. *Cell Metab*, 30(4):630-655.
- [89] Jadeja RN, Thounaojam MC, Bartoli M, Martin PM (2020). Implications of NAD(+) Metabolism in the Aging Retina and Retinal Degeneration. *Oxid Med Cell Longev*, 2020(2692794).
- [90] Chaturvedi P, Tyagi SC (2018). NAD(+): A big player in cardiac and skeletal muscle remodeling and aging. *J Cell Physiol*, 233(3):1895-1896.
- [91] Levine DC, Hong H, Weidemann BJ, Ramsey KM, Affinati AH, Schmidt M S, et al. (2020). NAD(+) Controls Circadian Reprogramming through PER2 Nuclear Translocation to Counter Aging. *Mol Cell*, 78(5):835-849.
- [92] Mitchell SJ, Bernier M, Aon MA, Cortassa S, Kim EY, Fang EF, et al. (2018). Nicotinamide Improves Aspects of Healthspan, but Not Lifespan, in Mice. *Cell Metab*, 27(3):667-676.
- [93] Zhang H, Ryu D, Wu Y, Gariani K, Wang X, Luan P, et al. (2016). NAD(+) repletion improves mitochondrial and stem cell function and enhances life span in mice. *Science*, 352(6292):1436-1443.
- [94] Lucena-Cacace A, Otero-Albiol D, Jimenez-Garcia M P, Munoz-Galvan S, Carnero A (2018). NAMPT Is a Potent Oncogene in Colon Cancer Progression that Modulates Cancer Stem Cell Properties and Resistance to Therapy through Sirt1 and PARP. *Clin Cancer Res*, 24(5):1202-1215.
- [95] Zhang H, Zhang N, Liu Y, Su P, Liang Y, Li Y, et al. (2019). Epigenetic Regulation of NAMPT by NAMPT-AS Drives Metastatic Progression in Triple-Negative Breast Cancer. *Cancer Res*, 79(13):3347-3359.
- [96] Ju H Q, Zhuang ZN, Li H, Tian T, Lu YX, Fan XQ, et al. (2016). Regulation of the Nampt-mediated NAD salvage pathway and its therapeutic implications in pancreatic cancer. *Cancer Lett*, 379(1):1-11.
- [97] Li X Q, Lei J, Mao LH, Wang Q L, Xu F, Ran T, et al. (2019). NAMPT and NAPRT, Key Enzymes in NAD Salvage Synthesis Pathway, Are of Negative Prognostic Value in Colorectal Cancer. *Front Oncol*, 9(736).
- [98] Chowdhry S, Zanca C, Rajkumar U, Koga T, Diao Y, Raviram R, et al. (2019). NAD metabolic dependency in cancer is shaped by gene amplification and enhancer remodelling. *Nature*, 569(7757):570-575.
- [99] Navas LE, Carnero A (2021). NAD(+) metabolism, stemness, the immune response, and cancer. *Signal Transduct Target Ther*, 6(1):2.
- [100] Han Y, Kim B, Cho U, Park I S, Kim SI, Dhanasekaran DN, et al. (2019). Mitochondrial fission causes cisplatin resistance under hypoxic conditions via ROS in ovarian cancer cells. *Oncogene*, 38(45):7089-7105.
- [101] Chang YJ, Chen KW, Chen L (2020). Mitochondrial ROS1 Increases Mitochondrial Fission and Respiration in Oral Squamous Cancer Carcinoma. *Cancers (Basel)*, 12(10):
- [102] Schopf B, Weissensteiner H, Schafer G, Fazzini F, Charoentong P, Naschberger A, et al. (2020). OXPHOS remodeling in high-grade prostate cancer involves mtDNA mutations and increased succinate oxidation. *Nat Commun*, 11(1):1487.
- [103] Li X, Guo X, Li D, Du X, Yin C, Chen C, et al. (2018). Multi-regional sequencing reveals intratumor heterogeneity and positive selection of somatic mtDNA mutations in hepatocellular carcinoma and colorectal cancer. *Int J Cancer*, 143(5):1143-1152.
- [104] Tseng LM, Yin PH, Chi CW, Hsu C Y, Wu CW, Lee LM, et al. (2006). Mitochondrial DNA mutations and mitochondrial DNA depletion in breast cancer. *Genes Chromosomes Cancer*, 45(7):629-638.
- [105] Srinivasan S, Guha M, Dong DW, Whelan KA, Ruthel G, Uchikado Y, et al. (2016). Disruption of cytochrome c oxidase function induces the Warburg effect and metabolic reprogramming. *Oncogene*, 35(12):1585-1595.
- [106] Kumar R, Bhat TA, Walsh EM, Chaudhary AK, O'Malley J, Rhim J , et al. (2019). Cytochrome c Deficiency Confers Apoptosome and Mitochondrial Dysfunction in African-American Men with Prostate Cancer. *Cancer Res*, 79(7):1353-1368.
- [107] Gomes AP, Ilter D, Low V, Endress JE, Fernandez-Garcia J, Rosenzweig A, et al. (2020). Age-induced accumulation of methylmalonic acid promotes tumour progression. *Nature*, 585(7824):283-287.
- [108] (2020). Serum Methylmalonic Acid Mediates Aging-Related Cancer Aggressiveness. *Cancer Discov*, 10(10):1441.
- [109] Jacobs KR, Castellano-Gonzalez G, Guillemin GJ, Lovejoy DB (2017). Major Developments in the Design of Inhibitors along the Kynurenine Pathway. *Curr Med Chem*, 24(23):2471-2495.
- [110] Liu CL, Cheng SP, Chen MJ, Lin CH, Chen SN, Kuo Y H, et al. (2020). Quinolate Phosphoribosyltransferase Promotes Invasiveness of Breast Cancer Through Myosin Light Chain Phosphorylation. *Front Endocrinol (Lausanne)*, 11(621944).
- [111] Ho P C, Bihuniak JD, Macintyre A N, Staron M, Liu X, Amezcua R, et al. (2015). Phosphoenolpyruvate

- Is a Metabolic Checkpoint of Anti-tumor T Cell Responses. *Cell*, 162(6):1217-1228.
- [112] Chin R M, Fu X, Pai M Y, Vergnes L, Hwang H, Deng G, et al. (2014). The metabolite alpha-ketoglutarate extends lifespan by inhibiting ATP synthase and TOR. *Nature*, 510(7505):397-401.
- [113] Su Y, Wang T, Wu N, Li D, Fan X, Xu Z, et al. (2019). Alpha-ketoglutarate extends Drosophila lifespan by inhibiting mTOR and activating AMPK. *Aging* (Albany NY), 11(12):4183-4197.
- [114] Asadi SA, Edgar D, Liao CY, Hsu YM, Lucanic M, Asadi SA, et al. (2020). Alpha-Ketoglutarate, an Endogenous Metabolite, Extends Lifespan and Compresses Morbidity in Aging Mice. *Cell Metab*, 32(3):447-456.
- [115] Zhang Z, He C, Gao Y, Zhang L, Song Y, Zhu T, et al. (2021). alpha-ketoglutarate delays age-related fertility decline in mammals. *Aging Cell*, 20(2): e13291.
- [116] Wang Y, Deng P, Liu Y, Wu Y, Chen Y, Guo Y, et al. (2020). Alpha-ketoglutarate ameliorates age-related osteoporosis via regulating histone methylations. *Nat Commun*, 11(1):5596.
- [117] Tseng CW, Kuo WH, Chan SH, Chan HL, Chang KJ, Wang LH (2018). Transketolase Regulates the Metabolic Switch to Control Breast Cancer Cell Metastasis via the alpha-Ketoglutarate Signaling Pathway. *Cancer Res*, 78(11):2799-2812.
- [118] Atlante S, Visintin A, Marini E, Savoia M, Dianzani C, Giorgis M, et al. (2018). alpha-ketoglutarate dehydrogenase inhibition counteracts breast cancer-associated lung metastasis. *Cell Death Dis*, 9(7):756.
- [119] Li S, Fu C, Zhao Y, He J (2019). Intervention with alpha-Ketoglutarate Ameliorates Colitis-Related Colorectal Carcinoma via Modulation of the Gut Microbiome. *Biomed Res Int*, 2019(8020785).
- [120] Wang X, Liu R, Qu X, Yu H, Chu H, Zhang Y, et al. (2019). alpha-Ketoglutarate-Activated NF-kappaB Signaling Promotes Compensatory Glucose Uptake and Brain Tumor Development. *Mol Cell*, 76(1):148-162.
- [121] Lamberts SW, van den Beld AW, van der Lely A J (1997). The endocrinology of aging. *Science*, 278(5337):419-424.
- [122] Veldhuis J D, Roemmich J N, Richmond E J, Bowers C Y (2006). Somatotropic and gonadotropic axes linkages in infancy, childhood, and the puberty-adult transition. *Endocr Rev*, 27(2):101-140.
- [123] Mazzoccoli G, Paziienza V, Piepoli A, Muscarella L A, Inglese M, De Cata A, et al. (2010). Hypothalamus-hypophysis-thyroid axis function in healthy aging. *J Biol Regul Homeost Agents*, 24(4):433-439.
- [124] Jansen S W, Akintola A A, Roelfsema F, van der Spoel E, Cobbaert C M, Ballieux B E, et al. (2015). Human longevity is characterised by high thyroid stimulating hormone secretion without altered energy metabolism. *Sci Rep*, 5(11525).
- [125] Zutinic A, Pijl H, Ballieux BE, Roelfsema F, Westendorp R, Blauw GJ, et al. (2020). Familial Longevity is Associated with an Attenuated Thyroidal Response to Recombinant Human Thyroid Stimulating Hormone. *J Clin Endocrinol Metab*, 105(7): e2572-e2580
- [126] Chu Y D, Yeh C T (2020). The Molecular Function and Clinical Role of Thyroid Stimulating Hormone Receptor in Cancer Cells. *Cells-Basel*, 9(7):
- [127] Lechler M C, Crawford E D, Groh N, Widmaier K, Jung R, Kirstein J, et al. (2017). Reduced Insulin/IGF-1 Signaling Restores the Dynamic Properties of Key Stress Granule Proteins during Aging. *Cell Rep*, 18(2):454-467.
- [128] Roitenberg N, Bejerano-Sagie M, Boucholez H, Moll L, Marques F C, Golodetzki L, et al. (2018). Modulation of caveolae by insulin/IGF-1 signaling regulates aging of *Caenorhabditis elegans*. *Embo Rep*, 19(8):
- [129] Gesing A, Wiesenborn D, Do A, Menon V, Schneider A, Victoria B, et al. (2017). A Long-lived Mouse Lacking Both Growth Hormone and Growth Hormone Receptor: A New Animal Model for Aging Studies. *J Gerontol A Biol Sci Med Sci*, 72(8):1054-1061.
- [130] Sun LY, Fang Y, Patki A, Koopman J J, Allison DB, Hill CM, et al. (2017). Longevity is impacted by growth hormone action during early postnatal period. *Elife*, 6(
- [131] Ben-Avraham D, Govindaraju D R, Budagov T, Fradin D, Durda P, Liu B, et al. (2017). The GH receptor exon 3 deletion is a marker of male-specific exceptional longevity associated with increased GH sensitivity and taller stature. *Sci Adv*, 3(6): e1602025.
- [132] Zhang G, Li J, Purkayastha S, Tang Y, Zhang H, Yin Y, et al. (2013). Hypothalamic programming of systemic ageing involving IKK-beta, NF-kappaB and GnRH. *Nature*, 497(7448):211-216.
- [133] Song YS, Kim M J, Sun HJ, Kim HH, Shin HS, Kim YA, et al. (2019). Aberrant Thyroid-Stimulating Hormone Receptor Signaling Increases VEGF-A and CXCL8 Secretion of Thyroid Cancer Cells, Contributing to Angiogenesis and Tumor Growth. *Clin Cancer Res*, 25(1):414-425.
- [134] Nieto H, Boelaert K (2016). Women In Cancer Thematic Review: Thyroid-stimulating hormone in thyroid cancer: does it matter? *Endocr Relat Cancer*, 23(11): T109-T121.
- [135] Chu YD, Yeh CT (2020). The Molecular Function and Clinical Role of Thyroid Stimulating Hormone Receptor in Cancer Cells. *Cells-Basel*, 9(7):
- [136] Long X, Xiong W, Zeng X, Qi L, Cai Y, et al. (2019). Cancer-associated fibroblasts promote cisplatin resistance in bladder cancer cells by increasing IGF-1/ERbeta/Bcl-2 signalling. *Cell Death Dis*, 10(5):375.
- [137] Sanchez-Lopez E, Flashner-Abramson E, Shalpour S, Zhong Z, Taniguchi K, Levitzki A, et al. (2016). Targeting colorectal cancer via its microenvironment by inhibiting IGF-1 receptor-insulin receptor substrate and STAT3 signaling. *Oncogene*, 35(20):2634-2644.
- [138] Shi H, Fang W, Liu M, Fu D (2017). Complement component 1, q subcomponent binding protein (C1QBP) in lipid rafts mediates hepatic metastasis of pancreatic cancer by regulating IGF-1/IGF-1R signaling. *Int J Cancer*, 141(7):1389-1401.

- [139] Muoio MG, Talia M, Lappano R, Sims AH, Vella V, Cirillo F, et al. (2021). Activation of the S100A7/RAGE Pathway by IGF-1 Contributes to Angiogenesis in Breast Cancer. *Cancers (Basel)*, 13(4):
- [140] Arumugam A, Subramani R, Nandy SB, Terreros D, Dwivedi A K, Saltzstein E, et al. (2019). Silencing growth hormone receptor inhibits estrogen receptor negative breast cancer through ATP-binding cassette sub-family G member 2. *Exp Mol Med*, 51(1):1-13.
- [141] Chhabra Y, Wong HY, Nikolajsen LF, Steinocher H, Papadopulos A, Tunny KA, et al. (2018). A growth hormone receptor SNP promotes lung cancer by impairment of SOCS2-mediated degradation. *Oncogene*, 37(4):489-501.
- [142] Evans A, Jamieson S M, Liu D X, Wilson W R, Perry J K (2016). Growth hormone receptor antagonism suppresses tumour regrowth after radiotherapy in an endometrial cancer xenograft model. *Cancer Lett*, 379(1):117-123.
- [143] Swerdlow AJ, Cooke R, Beckers D, Borgstrom B, Butler G, Carel JC, et al. (2017). Cancer Risks in Patients Treated with Growth Hormone in Childhood: The SAGhE European Cohort Study. *J Clin Endocrinol Metab*, 102(5):1661-1672.
- [144] Podlutzky A, Valcarcel-Ares M N, Yancey K, Podlutzkaya V, Nagykaladi E, Gautam T, et al. (2017). The GH/IGF-1 axis in a critical period early in life determines cellular DNA repair capacity by altering transcriptional regulation of DNA repair-related genes: implications for the developmental origins of cancer. *Geroscience*, 39(2):147-160.
- [145] Zhang J, Zhang J, Wei Y, Li Q, Wang Q (2019). ACTL6A regulates follicle-stimulating hormone-driven glycolysis in ovarian cancer cells via PGK1. *Cell Death Dis*, 10(11):811.
- [146] Aune D, Chan DS, Vieira A R, Navarro RD, Vieira R, Greenwood DC, et al. (2013). Red and processed meat intake and risk of colorectal adenomas: a systematic review and meta-analysis of epidemiological studies. *Cancer Causes Control*, 24(4):611-627.
- [147] Christensen B C, Houseman E A, Marsit C J, Zheng S, Wrensch M R, Wiemels J L, et al. (2009). Aging and environmental exposures alter tissue-specific DNA methylation dependent upon CpG island context. *Plos Genet*, 5(8): e1000602.
- [148] Bell JT, Tsai PC, Yang T P, Pidsley R, Nisbet J, Glass D, et al. (2012). Epigenome-wide scans identify differentially methylated regions for age and age-related phenotypes in a healthy ageing population. *Plos Genet*, 8(4): e1002629.
- [149] Horvath S (2013). DNA methylation age of human tissues and cell types. *Genome Biol*, 14(10): R115.
- [150] Horvath S, Pirazzini C, Bacalini MG, Gentilini D, Di Blasio AM, Delledonne M, et al. (2015). Decreased epigenetic age of PBMCs from Italian semi-supercentenarians and their offspring. *Aging (Albany NY)*, 7(12):1159-1170.
- [151] Lim U, Song M A (2018). DNA Methylation as a Biomarker of Aging in Epidemiologic Studies. *Methods Mol Biol*, 1856(219-231).
- [152] Horvath S, Pirazzini C, Bacalini MG, Gentilini D, Di Blasio AM, Delledonne M, et al. (2015). Decreased epigenetic age of PBMCs from Italian semi-supercentenarians and their offspring. *Aging (Albany NY)*, 7(12):1159-1170.
- [153] Armstrong NJ, Mather KA, Thalamuthu A, Wright MJ, Trollor JN, Ames D, et al. (2017). Aging, exceptional longevity and comparisons of the Hannum and Horvath epigenetic clocks. *Epigenomics-Uk*, 9(5):689-700.
- [154] Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sada S, et al. (2013). Genome-wide methylation profiles reveal quantitative views of human aging rates. *Mol Cell*, 49(2):359-367.
- [155] Horvath S (2013). DNA methylation age of human tissues and cell types. *Genome Biol*, 14(10): R115.
- [156] Levine M E, Hosgood H D, Chen B, Absher D, Assimes T, Horvath S (2015). DNA methylation age of blood predicts future onset of lung cancer in the women's health initiative. *Aging (Albany NY)*, 7(9):690-700.
- [157] Zheng Y, Joyce B T, Colicino E, Liu L, Zhang W, Dai Q, et al. (2016). Blood Epigenetic Age may Predict Cancer Incidence and Mortality. *Ebiomedicine*, 5(68-73).
- [158] Ambatipudi S, Horvath S, Perrier F, Cuenin C, Hernandez-Vargas H, Le Calvez-Kelm F, et al. (2017). DNA methylome analysis identifies accelerated epigenetic ageing associated with postmenopausal breast cancer susceptibility. *Eur J Cancer*, 75(299-307).
- [159] Dugue PA, Bassett JK, Wong EM, Joo JE, Li S, Yu C, et al. (2021). Biological Aging Measures Based on Blood DNA Methylation and Risk of Cancer: A Prospective Study. *JNCI Cancer Spectr*, 5(1):a109.
- [160] Chen X, Zhang J, Ruan W, Huang M, Wang C, Wang H, et al. (2020). Urine DNA methylation assay enables early detection and recurrence monitoring for bladder cancer. *J Clin Invest*, 130(12):6278-6289.
- [161] Liu B, Ricarte F J, Mallisetty A, Villani C, Kottorou A, Rodgers K, et al. (2020). Detection of Promoter DNA Methylation in Urine and Plasma Aids the Detection of Non-Small Cell Lung Cancer. *Clin Cancer Res*, 26(16):4339-4348.
- [162] Zhang M, Wang Y, Wang Y, Jiang L, Li X, Gao H, et al. (2020). Integrative Analysis of DNA Methylation and Gene Expression to Determine Specific Diagnostic Biomarkers and Prognostic Biomarkers of Breast Cancer. *Front Cell Dev Biol*, 8(529386).
- [163] Wu L, Yang Y, Guo X, Shu X O, Cai Q, Shu X, et al. (2020). An integrative multi-omics analysis to identify candidate DNA methylation biomarkers related to prostate cancer risk. *Nat Commun*, 11(1):3905.
- [164] Issa JP, Vertino PM, Boehm CD, Newsham IF, Baylin SB (1996). Switch from monoallelic to biallelic human IGF2 promoter methylation during aging and carcinogenesis. *Proc Natl Acad Sci U S A*, 93(21):11757-11762.

- [165] Fu VX, Dobosy JR, Desotelle JA, Almassi N, Ewald JA, Srinivasan R, et al. (2008). Aging and cancer-related loss of insulin-like growth factor 2 imprinting in the mouse and human prostate. *Cancer Res*, 68(16):6797-6802.
- [166] Ratajczak MZ, Kucia M, Liu R, Shin DM, Bryndza E, Masternak MM, et al. (2011). RasGrf1: genomic imprinting, VSEs, and aging. *Aging (Albany NY)*, 3(7):692-697.
- [167] Takamaru H, Yamamoto E, Suzuki H, Nojima M, Maruyama R, Yamano HO, et al. (2012). Aberrant methylation of RASGRF1 is associated with an epigenetic field defect and increased risk of gastric cancer. *Cancer Prev Res (Phila)*, 5(10):1203-1212.
- [168] Jiang N, Du G, Tobias E, Wood JG, Whitaker R, Neretti N, et al. (2013). Dietary and genetic effects on age-related loss of gene silencing reveal epigenetic plasticity of chromatin repression during aging. *Aging (Albany NY)*, 5(11):813-824.
- [169] Carone DM, Lawrence JB (2013). Heterochromatin instability in cancer: from the Barr body to satellites and the nuclear periphery. *Semin Cancer Biol*, 23(2):99-108.
- [170] Chen K, Wang S, Sun QW, Zhang B, Ullah M, Sun Z (2021). Klotho Deficiency Causes Heart Aging via Impairing the Nrf2-GR Pathway. *Circ Res*, 128(4):492-507.
- [171] Xu X, Hao Y, Zhong Q, Hang J, Zhao Y, Qiao J (2020). Low KLOTHO level related to aging is associated with diminished ovarian reserve. *Fertil Steril*, 114(6):1250-1255.
- [172] Ullah M, Sun Z (2019). Klotho Deficiency Accelerates Stem Cells Aging by Impairing Telomerase Activity. *J Gerontol A Biol Sci Med Sci*, 74(9):1396-1407.
- [173] Leon J, Moreno AJ, Garay BI, Chalkley RJ, Burlingame A L, Wang D, et al. (2017). Peripheral Elevation of a Klotho Fragment Enhances Brain Function and Resilience in Young, Aging, and alpha-Synuclein Transgenic Mice. *Cell Rep*, 20(6):1360-1371.
- [174] Shmulevich R, Nissim TB, Wolf I, Merenbakh-Lamin K, Fishman D, Sekler I, et al. (2020). Klotho rewires cellular metabolism of breast cancer cells through alteration of calcium shuttling and mitochondrial activity. *Oncogene*, 39(24):4636-4649.
- [175] Arbel R T, Shahmoon S, Zigmond E, Etan T, Merenbakh-Lamin K, Pasmanik-Chor M, et al. (2019). Klotho suppresses colorectal cancer through modulation of the unfolded protein response. *Oncogene*, 38(6):794-807.
- [176] Liu Y, Pan J, Pan X, Wu L, Bian J, Lin Z, et al. (2019). Klotho-mediated targeting of CCL2 suppresses the induction of colorectal cancer progression by stromal cell senescent microenvironments. *Mol Oncol*, 13(11):2460-2475.
- [177] Zhou X, Fang X, Jiang Y, Geng L, Li X, Li Y, et al. (2017). Klotho, an anti-aging gene, acts as a tumor suppressor and inhibitor of IGF-1R signaling in diffuse large B cell lymphoma. *J Hematol Oncol*, 10(1):37.
- [178] Pont A R, Sadri N, Hsiao S J, Smith S, Schneider R J (2012). mRNA decay factor AUF1 maintains normal aging, telomere maintenance, and suppression of senescence by activation of telomerase transcription. *Mol Cell*, 47(1):5-15.
- [179] He J, Jiang F, Liang L, Wang DJ, Wei WX, Ji PP, et al. (2017). Targeting of AUF1 to vascular endothelial cells as a novel anti-aging therapy. *J Geriatr Cardiol*, 14(8):515-523.
- [180] Li X, Tian Z, Jin H, Xu J, Hua X, Yan H, et al. (2018). Decreased c-Myc mRNA Stability via the MicroRNA 141-3p/AUF1 Axis Is Crucial for p63alpha Inhibition of Cyclin D1 Gene Transcription and Bladder Cancer Cell Tumorigenicity. *Mol Cell Biol*, 38(21):
- [181] Zucconi BE, Wilson GM (2011). Modulation of neoplastic gene regulatory pathways by the RNA-binding factor AUF1. *Front Biosci (Landmark Ed)*, 16(2307-2325).
- [182] Tian XY, Li J, Liu T H, Li DN, Wang JJ, Zhang H, et al. (2020). The overexpression of AUF1 in colorectal cancer predicts a poor prognosis and promotes cancer progression by activating ERK and AKT pathways. *Cancer Med*, 9(22):8612-8623.
- [183] Yang Y, Kang P, Gao J, Xu C, Wang S, Jin H, et al. (2014). AU-binding factor 1 expression was correlated with metadherin expression and progression of hepatocellular carcinoma. *Tumour Biol*, 35(3):2747-2751.
- [184] Du X, Wang J M, Zhang D L, Wu T, Zeng X Y, Jiang J Y, et al. (2021). AUF1 Promotes Proliferation and Invasion of Thyroid Cancer via Downregulation of ZBTB2 and Subsequent TRIM58. *Front Oncol*, 11(681736).
- [185] Herskovits AZ, Hunter T A, Maxwell N, Pereira K, Whittaker CA, Valdez G, et al. (2018). SIRT1 deacetylase in aging-induced neuromuscular degeneration and amyotrophic lateral sclerosis. *Aging Cell*, 17(6): e12839.
- [186] Kong D, Zhan Y, Liu Z, Ding T, Li M, Yu H, et al. (2016). SIRT1-mediated ERbeta suppression in the endothelium contributes to vascular aging. *Aging Cell*, 15(6):1092-1102.
- [187] Zhang S, Yang Y, Huang S, Deng C, Zhou S, Yang J, et al. (2019). SIRT1 inhibits gastric cancer proliferation and metastasis via STAT3/MMP-13 signaling. *J Cell Physiol*, 234(9):15395-15406.
- [188] Shi L, Tang X, Qian M, Liu Z, Meng F, Fu L, et al. (2018). A SIRT1-centered circuitry regulates breast cancer stemness and metastasis. *Oncogene*, 37(49):6299-6315.
- [189] Cheng F, Su L, Yao C, Liu L, Shen J, Liu C, et al. (2016). SIRT1 promotes epithelial-mesenchymal transition and metastasis in colorectal cancer by regulating Fra-1 expression. *Cancer Lett*, 375(2):274-283.
- [190] Wang T W, Chern E, Hsu C W, Tseng K C, Chao H M (2020). SIRT1-Mediated Expression of CD24 and Epigenetic Suppression of Novel Tumor Suppressor miR-1185-1 Increases Colorectal Cancer Stemness. *Cancer Res*, 80(23):5257-5269.

- [191] So D, Shin HW, Kim J, Lee M, Myeong J, Chun YS, et al. (2018). Cervical cancer is addicted to SIRT1 disarming the AIM2 antiviral defense. *Oncogene*, 37(38):5191-5204.
- [192] Zindy F, Quelle DE, Roussel MF, Sherr CJ (1997). Expression of the p16INK4a tumor suppressor versus other INK4 family members during mouse development and aging. *Oncogene*, 15(2):203-211.
- [193] Chen H, Gu X, Liu Y, Wang J, Wirt S E, Bottino R, et al. (2011). PDGF signalling controls age-dependent proliferation in pancreatic beta-cells. *Nature*, 478(7369):349-355.
- [194] Berent-Maoz B, Montecino-Rodriguez E, Signer RA, Dorshkind K (2012). Fibroblast growth factor-7 partially reverses murine thymocyte progenitor aging by repression of Ink4a. *Blood*, 119(24):5715-5721.
- [195] Baker D J, Wijshake T, Tchkonja T, LeBrasseur NK, Childs B G, van de Sluis B, et al. (2011). Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature*, 479(7372):232-236.
- [196] Zhu P, Zhang C, Gao Y, Wu F, Zhou Y, Wu WS (2019). The transcription factor Slug represses p16(Ink4a) and regulates murine muscle stem cell aging. *Nat Commun*, 10(1):2568.
- [197] Du W, Liu Z, Zhu W, Li T, Zhu Z, Wei L, et al. (2019). CSN6 promotes tumorigenesis of gastric cancer by ubiquitin-independent proteasomal degradation of p16(INK4a). *Cancer Biol Med*, 16(3):514-529.
- [198] Su W, Hong L, Xu X, Huang S, Herpai D, Li L, et al. (2018). miR-30 disrupts senescence and promotes cancer by targeting both p16(INK4A) and DNA damage pathways. *Oncogene*, 37(42):5618-5632.
- [199] Liu Y, Johnson S M, Fedorow Y, Rogers A B, Yuan H, Krishnamurthy J, et al. (2011). Expression of p16(INK4a) prevents cancer and promotes aging in lymphocytes. *Blood*, 117(12):3257-3267.
- [200] Qin X, Du D, Chen Q, Wu M, Wu T, Wen J, et al. (2019). Metformin prevents murine ovarian aging. *Aging (Albany NY)*, 11(11):3785-3794.
- [201] Zhao P, Sui B D, Liu N, Lv Y J, Zheng C X, Lu Y B, et al. (2017). Anti-aging pharmacology in cutaneous wound healing: effects of metformin, resveratrol, and rapamycin by local application. *Aging Cell*, 16(5):1083-1093.
- [202] Cai H, Han B, Hu Y, Zhao X, He Z, Chen X, et al. (2020). Metformin attenuates the Dgalactose-induced aging process via the UPR through the AMPK/ERK1/2 signaling pathways. *Int J Mol Med*, 45(3):715-730.
- [203] Fatemi I, Khaluoi A, Kaeidi A, Shamsizadeh A, Heydari S, Allahtavakoli M A (2018). Protective effect of metformin on D-galactose-induced aging model in mice. *Iran J Basic Med Sci*, 21(1):19-25.
- [204] Ahmadi S, Razazan A, Nagpal R, Jain S, Wang B, Mishra SP, et al. (2020). Metformin Reduces Aging-Related Leaky Gut and Improves Cognitive Function by Beneficially Modulating Gut Microbiome/Goblet Cell/Mucin Axis. *J Gerontol A Biol Sci Med Sci*, 75(7): e9-e21.
- [205] Anisimov VN, Piskunova TS, Popovich IG, Zabezhinski MA, Tyndyk ML, Egormin PA, et al. (2010). Gender differences in metformin effect on aging, life span and spontaneous tumorigenesis in 129/Sv mice. *Aging (Albany NY)*, 2(12):945-958.
- [206] Anisimov V N, Popovich I G, Zabezhinski M A, Egormin P A, Yurova M N, Semenchenko A V, et al. (2015). Sex differences in aging, life span and spontaneous tumorigenesis in 129/Sv mice neonatally exposed to metformin. *Cell Cycle*, 14(1):46-55.
- [207] Zhu X, Shen W, Liu Z, Sheng S, Xiong W, He R, et al. (2020). Effect of Metformin on Cardiac Metabolism and Longevity in Aged Female Mice. *Front Cell Dev Biol*, 8(626011).
- [208] Fang J, Yang J, Wu X, Zhang G, Li T, Wang X, et al. (2018). Metformin alleviates human cellular aging by upregulating the endoplasmic reticulum glutathione peroxidase 7. *Aging Cell*, 17(4):e12765.
- [209] Fitzgerald JC, Zimprich A, Carvajal BD, Schindler K M, Maurer B, Schulte C, et al. (2017). Metformin reverses TRAP1 mutation-associated alterations in mitochondrial function in Parkinson's disease. *Brain*, 140(9):2444-2459.
- [210] Sumantri S, Setiati S, Purnamasari D, Dewiasty E (2014). Relationship between metformin and frailty syndrome in elderly people with type 2 diabetes. *Acta Med Indones*, 46(3):183-188.
- [211] Samaras K, Makkar S, Crawford JD, Kochan NA, Wen W, Draper B, et al. (2020). Metformin Use Is Associated With Slowed Cognitive Decline and Reduced Incident Dementia in Older Adults With Type 2 Diabetes: The Sydney Memory and Ageing Study. *Diabetes Care*, 43(11):2691-2701.
- [212] Xia C, Liu C, He Z, Cai Y, Chen J (2020). Metformin inhibits cervical cancer cell proliferation by modulating PI3K/Akt-induced major histocompatibility complex class I-related chain A gene expression. *J Exp Clin Cancer Res*, 39(1):127.
- [213] Amable G, Martínez León E, Picco M E, Nemirovsky S I, Rozengurt E, Rey O (2020). Metformin inhibition of colorectal cancer cell migration is associated with rebuilt adherens junctions and FAK downregulation. *J Cell Physiol*, 235(11):8334-8344.
- [214] Hsieh L S, Liu S T, Chang Y L, Ho C L, Huang S M (2018). Metformin causes cancer cell death through downregulation of p53-dependent differentiated embryo chondrocyte 1. *J Biomed Sci*, 25(1):81.
- [215] Chen K, Qian W, Jiang Z, Cheng L, Li J, Sun L, et al. (2017). Metformin suppresses cancer initiation and progression in genetic mouse models of pancreatic cancer. *Mol Cancer*, 16(1):131.
- [216] Wang J C, Li G Y, Wang B, Han S X, Sun X, Jiang Y N, et al. (2019). Metformin inhibits metastatic breast cancer progression and improves chemosensitivity by inducing vessel normalization via PDGF-B downregulation. *J Exp Clin Cancer Res*, 38(1):235.
- [217] Sonnenblick A, Agbor-Tarh D, Bradbury I, Di Cosimo S, Azim H J, Fumagalli D, et al. (2017). Impact of Diabetes, Insulin, and Metformin Use on the Outcome of Patients With Human Epidermal Growth Factor

- Receptor 2-Positive Primary Breast Cancer: Analysis From the ALTO Phase III Randomized Trial. *J Clin Oncol*, 35(13):1421-1429.
- [218] Brown J R, Chan D K, Shank J J, Griffith K A, Fan H, Szulawski R, et al. (2020). Phase II clinical trial of metformin as a cancer stem cell-targeting agent in ovarian cancer. *JCI Insight*, 5(11):
- [219] Marrone K A, Zhou X, Forde P M, Purtell M, Brahmer J R, Hann C L, et al. (2018). A Randomized Phase II Study of Metformin plus Paclitaxel/Carboplatin/Bevacizumab in Patients with Chemotherapy-Naive Advanced or Metastatic Nonsquamous Non-Small Cell Lung Cancer. *Oncologist*, 23(7):859-865.
- [220] Kordes S, Pollak M N, Zwinderman A H, Mathot R A, Weterman M J, Beeker A, et al. (2015). Metformin in patients with advanced pancreatic cancer: a double-blind, randomised, placebo-controlled phase 2 trial. *Lancet Oncol*, 16(7):839-847.
- [221] Anisimov V N, Berstein L M, Popovich I G, Zabezhinski M A, Egorin P A, Piskunova T S, et al. (2011). If started early in life, metformin treatment increases life span and postpones tumors in female SHR mice. *Aging (Albany NY)*, 3(2):148-157.
- [222] Aguilar-Alonso P, Vera-Lopez O, Brambila-Colombes E, Segura-Badilla O, Avalos-Lopez R, Lazzano-Hernandez M, et al. (2018). Evaluation of Oxidative Stress in Cardiomyocytes during the Aging Process in Rats Treated with Resveratrol. *Oxid Med Cell Longev*, 2018(1390483).
- [223] Navarro-Cruz AR, Ramirez YAR, Ochoa-Velasco C, Brambila E, Avila-Sosa R, Perez-Fernandez S, et al. (2017). Effect of Chronic Administration of Resveratrol on Cognitive Performance during Aging Process in Rats. *Oxid Med Cell Longev*, 2017(8510761).
- [224] Kim E N, Lim J H, Kim M Y, Ban T H, Jang I A, Yoon H E, et al. (2018). Resveratrol, an Nrf2 activator, ameliorates aging-related progressive renal injury. *Aging (Albany NY)*, 10(1):83-99.
- [225] Kim E N, Kim M Y, Lim J H, Kim Y, Shin S J, Park C W, et al. (2018). The protective effect of resveratrol on vascular aging by modulation of the renin-angiotensin system. *Atherosclerosis*, 270(123-131).
- [226] Gerhardt E, Graber S, Szego E M, Moiso N, Martins L M, Outeiro T F, et al. (2011). Idebenone and resveratrol extend lifespan and improve motor function of HtrA2 knockout mice. *Plos One*, 6(12):e28855.
- [227] Mancuso R, Del V J, Modol L, Martinez A, Granado-Serrano A B, Ramirez-Nunez O, et al. (2014). Resveratrol improves motoneuron function and extends survival in SOD1(G93A) ALS mice. *Neurotherapeutics*, 11(2):419-432.
- [228] Porquet D, Casadesus G, Bayod S, Vicente A, Canudas A M, Vilaplana J, et al. (2013). Dietary resveratrol prevents Alzheimer's markers and increases life span in SAMP8. *Age (Dordr)*, 35(5):1851-1865.
- [229] Boo YC (2019). Human Skin Lightening Efficacy of Resveratrol and Its Analogs: From in Vitro Studies to Cosmetic Applications. *Antioxidants (Basel)*, 8(9):
- [230] Evans H M, Howe P R, Wong R H (2017). Effects of Resveratrol on Cognitive Performance, Mood and Cerebrovascular Function in Post-Menopausal Women; A 14-Week Randomised Placebo-Controlled Intervention Trial. *Nutrients*, 9(1):
- [231] Moran C, Scotto DPA, Bramham J, Moran A, Rooney B, De Vito G, et al. (2018). Effects of a Six-Month Multi-Ingredient Nutrition Supplement Intervention of Omega-3 Polyunsaturated Fatty Acids, vitamin D, Resveratrol, and Whey Protein on Cognitive Function in Older Adults: A Randomised, Double-Blind, Controlled Trial. *J Prev Alzheimers Dis*, 5(3):175-183.
- [232] Harper SA, Bassler JR, Peramsetty S, Yang Y, Roberts LM, Drummer D, et al. (2021). Resveratrol and exercise combined to treat functional limitations in late life: A pilot randomized controlled trial. *Exp Gerontol*, 143:111111.
- [233] Lin YS, Hsieh CY, Kuo TT, Lin CC, Lin C Y, Sher Y P (2020). Resveratrol-mediated ADAM9 degradation decreases cancer progression and provides synergistic effects in combination with chemotherapy. *Am J Cancer Res*, 10(11):3828-3837.
- [234] Kim J Y, Cho K H, Jeong B Y, Park C G, Lee H Y (2020). Zeb1 for RCP-induced oral cancer cell invasion and its suppression by resveratrol. *Exp Mol Med*, 52(7):1152-1163.
- [235] Alayev A, Salamon RS, Schwartz NS, Berman AY, Wiener SL, Holz MK (2017). Combination of Rapamycin and Resveratrol for Treatment of Bladder Cancer. *J Cell Physiol*, 232(2):436-446.
- [236] Singh CK, Chhabra G, Ndiaye MA, Siddiqui IA, Panackal JE, Mintie C A, et al. (2020). Quercetin-Resveratrol Combination for Prostate Cancer Management in TRAMP Mice. *Cancers (Basel)*, 12(8):
- [237] Hu WH, Chan GK, Duan R, Wang HY, Kong XP, Dong TT, et al. (2019). Synergy of Ginkgetin and Resveratrol in Suppressing VEGF-Induced Angiogenesis: A Therapy in Treating Colorectal Cancer. *Cancers (Basel)*, 11(12):
- [238] Wang H, Jia R, Lv T, Wang M, He S, Zhang X (2020). Resveratrol Suppresses Tumor Progression via Inhibiting STAT3/HIF-1alpha/VEGF Pathway in an Orthotopic Rat Model of Non-Small-Cell Lung Cancer (NSCLC). *Onco Targets Ther*, 13(7057-7063).
- [239] Zhou C, Qian W, Ma J, Cheng L, Jiang Z, Yan B, et al. (2019). Resveratrol enhances the chemotherapeutic response and reverses the stemness induced by gemcitabine in pancreatic cancer cells via targeting SREBP1. *Cell Prolif*, 52(1):e12514.
- [240] Dou X, Sun Y, Li J, Zhang J, Hao D, Liu W, et al. (2017). Short-term rapamycin treatment increases ovarian lifespan in young and middle-aged female mice. *Aging Cell*, 16(4):825-836.
- [241] An JY, Kerns KA, Ouellette A, Robinson L, Morris H D, Kaczorowski C, et al. (2020). Rapamycin rejuvenates oral health in aging mice. *Elife*, 9(

- [242] Wang T, Tsui B, Kreisberg J F, Robertson N A, Gross A M, Yu M K, et al. (2017). Epigenetic aging signatures in mice livers are slowed by dwarfism, calorie restriction and rapamycin treatment. *Genome Biol*, 18(1):57.
- [243] Bitto A, Ito TK, Pineda VV, LeTexier NJ, Huang HZ, Sutlief E, et al. (2016). Transient rapamycin treatment can increase lifespan and healthspan in middle-aged mice. *Elife*, 5:e16351
- [244] Strong R, Miller R A, Bogue M, Fernandez E, Javors M A, Libert S, et al. (2020). Rapamycin-mediated mouse lifespan extension: Late-life dosage regimes with sex-specific effects. *Aging Cell*, 19(11):e13269.
- [245] Birkisdottir M B, Jaarsma D, Brandt R, Barnhoorn S, van Vliet N, Imholz S, et al. (2021). Unlike dietary restriction, rapamycin fails to extend lifespan and reduce transcription stress in progeroid DNA repair-deficient mice. *Aging Cell*, 20(2):e13302.
- [246] Blagosklonny M V (2021). DNA- and telomere-damage does not limit lifespan: evidence from rapamycin. *Aging (Albany, NY.)*, 13(3):3167-3175.
- [247] Chung C L, Lawrence I, Hoffman M, Elgindi D, Nadhan K, Potnis M, et al. (2019). Topical rapamycin reduces markers of senescence and aging in human skin: an exploratory, prospective, randomized trial. *Geroscience*, 41(6):861-869.
- [248] Shen Y Q, Guerra-Librero A, Fernandez-Gil B I, Florido J, Garcia-Lopez S, Martinez-Ruiz L, et al. (2018). Combination of melatonin and rapamycin for head and neck cancer therapy: Suppression of AKT/mTOR pathway activation, and activation of mitophagy and apoptosis via mitochondrial function regulation. *J Pineal Res*, 64(3):
- [249] Imrali A, Mao X, Yeste-Velasco M, Shamash J, Lu Y (2016). Rapamycin inhibits prostate cancer cell growth through cyclin D1 and enhances the cytotoxic efficacy of cisplatin. *Am J Cancer Res*, 6(8):1772-1784.
- [250] Wu Q, Kiguchi K, Kawamoto T, Ajiki T, Traag J, Carbajal S, et al. (2007). Therapeutic effect of rapamycin on gallbladder cancer in a transgenic mouse model. *Cancer Res*, 67(8):3794-3800.
- [251] Buijssen J, van den Bogaard J, Jutten B, Belgers E, Sosef M, Leijtens J W, et al. (2015). A phase I-II study on the combination of rapamycin and short course radiotherapy in rectal cancer. *Radiother Oncol*, 116(2):214-220.
- [252] Day T A, Shirai K, O'Brien PE, Matheus MG, Godwin K, Sood AJ, et al. (2019). Inhibition of mTOR Signaling and Clinical Activity of Rapamycin in Head and Neck Cancer in a Window of Opportunity Trial. *Clin Cancer Res*, 25(4):1156-1164.
- [253] Ji N, Mukherjee N, Reyes RM, Gelfond J, Javors M, Meeks J J, et al. (2021). Rapamycin enhances BCG-specific gammadelta T cells during intravesical BCG therapy for non-muscle invasive bladder cancer: a randomized, double-blind study. *J Immunother Cancer*, 9(3):
- [254] Dantal J, Morelon E, Rostaing L, Goffin E, Brocard A, Tromme I, et al. (2018). Sirolimus for Secondary Prevention of Skin Cancer in Kidney Transplant Recipients: 5-Year Results. *J Clin Oncol*, 36(25):2612-2620.
- [255] Anisimov VN, Zabezhinski MA, Popovich IG, Piskunova TS, Semenchenko A V, Tyndyk M L, et al. (2010). Rapamycin extends maximal lifespan in cancer-prone mice. *Am J Pathol*, 176(5):2092-2097.
- [256] Wilkinson JE, Burmeister L, Brooks SV, Chan CC, Friedline S, Harrison DE, et al. (2012). Rapamycin slows aging in mice. *Aging Cell*, 11(4):675-682.
- [257] Kapahi P, Kaeberlein M, Hansen M (2017). Dietary restriction and lifespan: Lessons from invertebrate models. *Ageing Res Rev*, 39(3-14).
- [258] Simsek B, Yanar K, Kansu A D, Belce A, Aydin S, Cakatay U (2019). Caloric restriction improves the redox homeostasis in the aging male rat heart even when started in middle-adulthood and when the body weight is stable. *Biogerontology*, 20(1):127-140.
- [259] Gutierrez-Casado E, Khraiweh H, Lopez-Dominguez J A, Montero-Guisado J, Lopez-Lluch G, Navas P, et al. (2019). The Impact of Aging, Calorie Restriction and Dietary Fat on Autophagy Markers and Mitochondrial Ultrastructure and Dynamics in Mouse Skeletal Muscle. *J Gerontol A Biol Sci Med Sci*, 74(6):760-769.
- [260] Salvatore M F, Terrebonne J, Cantu MA, McInnis T R, Venable K, Kelley P, et al. (2017). Dissociation of Striatal Dopamine and Tyrosine Hydroxylase Expression from Aging-Related Motor Decline: Evidence from Calorie Restriction Intervention. *J Gerontol A Biol Sci Med Sci*, 73(1):11-20.
- [261] Sheng Y, Xia F, Chen L, Lv Y, Lv S, Yu J, et al. (2021). Differential Responses of White Adipose Tissue and Brown Adipose Tissue to Calorie Restriction During Aging. *J Gerontol A Biol Sci Med Sci*, 76(3):393-399.
- [262] Sziraki A, Tyshkovskiy A, Gladyshev V N (2018). Global remodeling of the mouse DNA methylome during aging and in response to calorie restriction. *Aging Cell*, 17(3):e12738.
- [263] Wang T, Tsui B, Kreisberg J F, Robertson NA, Gross AM, Yu MK, et al. (2017). Epigenetic aging signatures in mice livers are slowed by dwarfism, calorie restriction and rapamycin treatment. *Genome Biol*, 18(1):57.
- [264] Wolf NS, Penn PE, Jiang D, Fei RG, Pendergrass W R (1995). Caloric restriction: conservation of in vivo cellular replicative capacity accompanies life-span extension in mice. *Exp Cell Res*, 217(2):317-323.
- [265] Zha Y, Taguchi T, Nazneen A, Shimokawa I, Higami Y, Razzaque M S (2008). Genetic suppression of GH-IGF-1 activity, combined with lifelong caloric restriction, prevents age-related renal damage and prolongs the life span in rats. *Am J Nephrol*, 28(5):755-764.
- [266] Mitterberger MC, Mattesich M, Zwerschke W (2014). Bariatric surgery and diet-induced long-term caloric restriction protect subcutaneous adipose-derived stromal/progenitor cells and prolong their life span in formerly obese humans. *Exp Gerontol*, 56(106-113).

- [267] Willcox BJ, Willcox DC, Todoriki H, Fujiyoshi A, Yano K, He Q, et al. (2007). Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. *Ann N Y Acad Sci*, 1114(434-455).
- [268] Simone BA, Palagani A, Strickland K, Ko K, Jin L, Lim MK, et al. (2018). Caloric restriction counteracts chemotherapy-induced inflammation and increases response to therapy in a triple negative breast cancer model. *Cell Cycle*, 17(13):1536-1544.
- [269] Rogozina OP, Bonorden MJ, Seppanen CN, Grande JP, Cleary M P (2011). Effect of chronic and intermittent calorie restriction on serum adiponectin and leptin and mammary tumorigenesis. *Cancer Prev Res (Phila)*, 4(4):568-581.
- [270] Alidadi M, Banach M, Guest PC, Bo S, Jamialahmadi T, Sahebkar A (2020). The effect of caloric restriction and fasting on cancer. *Semin Cancer Biol*,
- [271] Lope V, Martin M, Castello A, Ruiz A, Casas A M, Baena-Canada J M, et al. (2019). Overeating, caloric restriction and breast cancer risk by pathologic subtype: the EPIGEICAM study. *Sci Rep*, 9(1):3904.
- [272] Weindruch R, Walford R L (1982). Dietary restriction in mice beginning at 1 year of age: effect on life-span and spontaneous cancer incidence. *Science*, 215(4538):1415-1418.
- [273] Lee C K, Pugh T D, Klopp R G, Edwards J, Allison D B, Weindruch R, et al. (2004). The impact of alpha-lipoic acid, coenzyme Q10 and caloric restriction on life span and gene expression patterns in mice. *Free Radic Biol Med*, 36(8):1043-1057.
- [274] Newman J C, Covarrubias A J, Zhao M, Yu X, Gut P, Ng C P, et al. (2017). Ketogenic Diet Reduces Midlife Mortality and Improves Memory in Aging Mice. *Cell Metab*, 26(3):547-557.
- [275] Yu Y, Wang F, Wang J, Zhang D, Zhao X (2020). Ketogenic diet attenuates aging-associated myocardial remodeling and dysfunction in mice. *Exp Gerontol*, 140(111058).
- [276] Wallace M A, Aguirre N W, Marcotte G R, Marshall A G, Baehr L M, Hughes D C, et al. (2021). The ketogenic diet preserves skeletal muscle with aging in mice. *Aging Cell*, 20(4):e13322.
- [277] Roberts M N, Wallace M A, Tomilov A A, Zhou Z, Marcotte G R, Tran D, et al. (2017). A Ketogenic Diet Extends Longevity and Healthspan in Adult Mice. *Cell Metab*, 26(3):539-546.
- [278] Rusek M, Pluta R, Ulamek-Kozioł M, Czuczwar S J (2019). Ketogenic Diet in Alzheimer's Disease. *Int J Mol Sci*, 20(16):
- [279] Zou Y, Fineberg S, Pearlman A, Feinman R D, Fine E J (2020). The effect of a ketogenic diet and synergy with rapamycin in a mouse model of breast cancer. *Plos One*, 15(12):e233662.
- [280] Zhang N, Liu C, Jin L, Zhang R, Wang T, Wang Q, et al. (2020). Ketogenic Diet Elicits Antitumor Properties through Inducing Oxidative Stress, Inhibiting MMP-9 Expression, and Rebalancing M1/M2 Tumor-Associated Macrophage Phenotype in a Mouse Model of Colon Cancer. *J Agric Food Chem*, 68(40):11182-11196.
- [281] Zahra A, Fath MA, Opat E, Mapuskar KA, Bhatia SK, Ma DC, et al. (2017). Consuming a Ketogenic Diet while Receiving Radiation and Chemotherapy for Locally Advanced Lung Cancer and Pancreatic Cancer: The University of Iowa Experience of Two Phase 1 Clinical Trials. *Radiat Res*, 187(6):743-754.
- [282] Mukherjee P, Augur ZM, Li M, Hill C, Greenwood B, Domin M A, et al. (2019). Therapeutic benefit of combining calorie-restricted ketogenic diet and glutamine targeting in late-stage experimental glioblastoma. *Commun Biol*, 2(200).
- [283] Khodabakhshi A, Akbari M E, Mirzaei H R, Mehrad-Majd H, Kalamian M, D avoodi S H (2020). Feasibility, Safety, and Beneficial Effects of MCT-Based Ketogenic Diet for Breast Cancer Treatment: A Randomized Controlled Trial Study. *Nutr Cancer*, 72(4):627-634.
- [284] Hagihara K, Kajimoto K, Osaga S, Nagai N, Shimosegawa E, Nakata H, et al. (2020). Promising Effect of a New Ketogenic Diet Regimen in Patients with Advanced Cancer. *Nutrients*, 12(5):
- [285] Iyikesici M S (2019). Feasibility study of metabolically supported chemotherapy with weekly carboplatin/paclitaxel combined with ketogenic diet, hyperthermia and hyperbaric oxygen therapy in metastatic non-small cell lung cancer. *Int J Hyperthermia*, 36(1):446-455.
- [286] Yang X, Yu D, Xue L, Li H, Du J (2020). Probiotics modulate the microbiota-gut-brain axis and improve memory deficits in aged SAMP8 mice. *Acta Pharm Sin B*, 10(3):475-487.
- [287] Hunsche C, Cruces J, De la Fuente M (2019). Improvement of Redox State and Functions of Immune Cells as Well as of Behavioral Response in Aged Mice After Two-Week Supplementation of Fermented Milk with Probiotics. *Curr Microbiol*, 76(11):1278-1289.
- [288] Collins FL, Rios-Arce ND, Schepper JD, Parameswaran N, McCabe LR (2017). The Potential of Probiotics as a Therapy for Osteoporosis. *Microbiol Spectr*, 5(4):
- [289] Lee HJ, Jeong JJ, Han MJ, Kim DH (2018). *Lactobacillus plantarum* C29 Alleviates TNBS-Induced Memory Impairment in Mice. *J Microbiol Biotechnol*, 28(1):175-179.
- [290] Sharma D, Kober MM, Bowe WP (2016). Anti-Aging Effects of Probiotics. *J Drugs Dermatol*, 15(1):9-12.
- [291] Akbari E, Asemi Z, Daneshvar K R, Bahmani F, Kouchaki E, Tamtaji O R, et al. (2016). Effect of Probiotic Supplementation on Cognitive Function and Metabolic Status in Alzheimer's Disease: A Randomized, Double-Blind and Controlled Trial. *Front Aging Neurosci*, 8(256).
- [292] Tamtaji O R, Heidari-Soureshjani R, Mirhosseini N, Kouchaki E, Bahmani F, Aghadavod E, et al. (2019). Probiotic and selenium co-supplementation, and the effects on clinical, metabolic and genetic status in

- Alzheimer's disease: A randomized, double-blind, controlled trial. *Clin Nutr*, 38(6):2569-2575.
- [293] Romo-Araiza A, Gutiérrez-Salmeán G, Galván E J, Hernández-Frausto M, Herrera-López G, Romo-Parra H, et al. (2018). Probiotics and Prebiotics as a Therapeutic Strategy to Improve Memory in a Model of Middle-Aged Rats. *Front Aging Neurosci*, 10(2018):1-11.
- [294] Liu Y, Zhao Y, Yang Y, Wang Z (2019). Effects of Probiotics, Prebiotics, and Synbiotics on Calcium Homeostasis and Bone Health With Aging: A Systematic Review. *Worldviews Evid Based Nurs*, 16(6):478-484.
- [295] Eslami M, Yousefi B, Kokhaei P, Hemati M, Nejad Z R, Arabkari V, et al. (2019). Importance of probiotics in the prevention and treatment of colorectal cancer. *J Cell Physiol*, 234(10):17127-17143.
- [296] Mendoza L (2019). Potential effect of probiotics in the treatment of breast cancer. *Oncol Rev*, 13(2):422.
- [297] He L, Yang H, Tang J, Liu Z, Chen Y, Lu B, et al. (2019). Intestinal probiotics *E. coli* Nissle 1917 as a targeted vehicle for delivery of p53 and Tum-5 to solid tumors for cancer therapy. *J Biol Eng*, 13(58):1-11.
- [298] Kazmierczak-Siedlecka K, Daca A, Fic M, van de Wetering T, Folwarski M, Makarewicz W (2020). Therapeutic methods of gut microbiota modification in colorectal cancer management - fecal microbiota transplantation, prebiotics, probiotics, and synbiotics. *Gut Microbes*, 11(6):1518-1530.
- [299] Toi M, Hirota S, Tomotaki A, Sato N, Hozumi Y, Anan K, et al. (2013). Probiotic Beverage with Soy Isoflavone Consumption for Breast Cancer Prevention: A Case-control Study. *Curr Nutr Food Sci*, 9(3):194-200.
- [300] Takada K, Shimokawa M, Takamori S, Shimamatsu S, Hirai F, Tagawa T, et al. (2021). Clinical impact of probiotics on the efficacy of anti-PD-1 monotherapy in patients with nonsmall cell lung cancer: A multicenter retrospective survival analysis study with inverse probability of treatment weighting. *Int J Cancer*, 149(2):473-482.
- [301] Arai J, Niikura R, Hayakawa Y, Kawahara T, Honda T, Hasatani K, et al. (2021). Use of Antibiotics and Probiotics Reduces the Risk of Metachronous Gastric Cancer after Endoscopic Resection. *Biology (Basel)*, 10(6):1-11.
- [302] Barcena C, Valdes-Mas R, Mayoral P, Garabaya C, Durand S, Rodriguez F, et al. (2019). Healthspan and lifespan extension by fecal microbiota transplantation into progeroid mice. *Nat Med*, 25(8):1234-1242.
- [303] Yu F, Jiang R, Han W, Zhan G, Xu X, Jiang X, et al. (2020). Gut microbiota transplantation from db/db mice induces diabetes-like phenotypes and alterations in Hippo signaling in pseudo germ-free mice. *Aging (Albany NY)*, 12(23):24156-24167.
- [304] Wang Z, Hua W, Li C, Chang H, Liu R, Ni Y, et al. (2019). Protective Role of Fecal Microbiota Transplantation on Colitis and Colitis-Associated Colon Cancer in Mice Is Associated With Treg Cells. *Front Microbiol*, 10(2498):1-11.
- [305] Rosshart S P, Vassallo B G, Angeletti D, Hutchinson D S, Morgan A P, Takeda K, et al. (2017). Wild Mouse Gut Microbiota Promotes Host Fitness and Improves Disease Resistance. *Cell*, 171(5):1015-1028.
- [306] Routy B, Le Chatelier E, Derosa L, Duong C, Alou M T, Daillere R, et al. (2018). Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science*, 359(6371):91-97.
- [307] Gopalakrishnan V, Spencer C N, Nezi L, Reuben A, Andrews M C, Karpinets T V, et al. (2018). Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science*, 359(6371):97-103.
- [308] de Clercq N C, van den Ende T, Prodan A, Hemke R, Davids M, Pedersen H K, et al. (2021). Fecal Microbiota Transplantation from Overweight or Obese Donors in Cachectic Patients with Advanced Gastroesophageal Cancer: A Randomized, Double-blind, Placebo-Controlled, Phase II Study. *Clin Cancer Res*, 27(11):2711-2721.
- [309] Zhu Y, Carvey P M, Ling Z (2006). Age-related changes in glutathione and glutathione-related enzymes in rat brain. *Brain Res*, 1090(1):35-44.
- [310] Tong J, Fitzmaurice P S, Moszczynska A, Mattina K, Ang L C, Boileau I, et al. (2016). Do glutathione levels decline in aging human brain? *Free Radic Biol Med*, 93(110-117):110-117.
- [311] Madeo F, Carmona-Gutierrez D, Kepp O, Kroemer G (2018). Spermidine delays aging in humans. *Aging (Albany NY)*, 10(8):2209-2211.
- [312] Meynial-Denis D (2016). Glutamine metabolism in advanced age. *Nutr Rev*, 74(4):225-236.
- [313] Kiss T, Giles C B, Tarantini S, Yabluchanskiy A, Balasubramanian P, Gautam T, et al. (2019). Nicotinamide mononucleotide (NMN) supplementation promotes anti-aging miRNA expression profile in the aorta of aged mice, predicting epigenetic rejuvenation and anti-atherogenic effects. *Geroscience*, 41(4):419-439.
- [314] de Picciotto N E, Gano L B, Johnson L C, Martens C R, Sindler A L, Mills K F, et al. (2016). Nicotinamide mononucleotide supplementation reverses vascular dysfunction and oxidative stress with aging in mice. *Aging Cell*, 15(3):522-530.
- [315] Hosseini L, Farokhi-Sisakht F, Badalzadeh R, Khabbaz A, Mahmoudi J, Sadigh-Eteghad S (2019). Nicotinamide Mononucleotide and Melatonin Alleviate Aging-induced Cognitive Impairment via Modulation of Mitochondrial Function and Apoptosis in the Prefrontal Cortex and Hippocampus. *Neuroscience*, 423(29-37):29-37.
- [316] Tarantini S, Valcarcel-Ares MN, Toth P, Yabluchanskiy A, Tucek Z, Kiss T, et al. (2019). Nicotinamide mononucleotide (NMN) supplementation rescues cerebrovascular endothelial function and neurovascular coupling responses and improves cognitive function in aged mice. *Redox Biol*, 24(101192):1-11.
- [317] Kiss T, Nyul-Toth A, Balasubramanian P, Tarantini S, Ahire C, Yabluchanskiy A, et al. (2020). Nicotinamide

- mononucleotide (NMN) supplementation promotes neurovascular rejuvenation in aged mice: transcriptional footprint of SIRT1 activation, mitochondrial protection, anti-inflammatory, and anti-apoptotic effects. *Geroscience*, 42(2):527-546.
- [318] Song J, Li J, Yang F, Ning G, Zhen L, Wu L, et al. (2019). Nicotinamide mononucleotide promotes osteogenesis and reduces adipogenesis by regulating mesenchymal stromal cells via the SIRT1 pathway in aged bone marrow. *Cell Death Dis*, 10(5):336.
- [319] Mills K F, Yoshida S, Stein L R, Grozio A, Kubota S, Sasaki Y, et al. (2016). Long-Term Administration of Nicotinamide Mononucleotide Mitigates Age-Associated Physiological Decline in Mice. *Cell Metab*, 24(6):795-806.
- [320] Cui C, Qi J, Deng Q, Chen R, Zhai D, Yu J (2016). Nicotinamide Mononucleotide Adenylyl Transferase 2: A Promising Diagnostic and Therapeutic Target for Colorectal Cancer. *Biomed Res Int*, 2016(1804)137.
- [321] Grozio A, Mills K F, Yoshino J, Bruzzone S, Sociali G, Tokizane K, et al. (2019). Slc12a8 is a nicotinamide mononucleotide transporter. *Nat Metab*, 1(1):47-57.
- [322] Zong Z, Liu J, Wang N, Yang C, Wang Q, Zhang W, et al. (2021). Nicotinamide mononucleotide inhibits hepatic stellate cell activation to prevent liver fibrosis via promoting PGE2 degradation. *Free Radic Biol Med*, 162(571)-581.
- [323] Schwarz C, Horn N, Benson G, Wrachtrup C I, Wurdack K, Pechlaner R, et al. (2020). Spermidine intake is associated with cortical thickness and hippocampal volume in older adults. *Neuroimage*, 221(11)7132.
- [324] Schroeder S, Hofer S J, Zimmermann A, Pechlaner R, Dammbrueck C, Pendl T, et al. (2021). Dietary spermidine improves cognitive function. *Cell Rep*, 35(2):108985.
- [325] Minois N, Rockenfeller P, Smith T K, Carmona-Gutierrez D (2014). Spermidine feeding decreases age-related locomotor activity loss and induces changes in lipid composition. *Plos One*, 9(7):e102435.
- [326] Wang J, Li S, Wang J, Wu F, Chen Y, Zhang H, et al. (2020). Spermidine alleviates cardiac aging by improving mitochondrial biogenesis and function. *Aging (Albany NY)*, 12(1):650-671.
- [327] Eisenberg T, Abdellatif M, Schroeder S, Primessnig U, Stekovic S, Pendl T, et al. (2016). Cardioprotection and lifespan extension by the natural polyamine spermidine. *Nat Med*, 22(12):1428-1438.
- [328] Fan J, Yang X, Li J, Shu Z, Dai J, Liu X, et al. (2017). Spermidine coupled with exercise rescues skeletal muscle atrophy from D-gal-induced aging rats through enhanced autophagy and reduced apoptosis via AMPK-FOXO3a signal pathway. *Oncotarget*, 8(11):17475-17490.
- [329] LaRocca T J, Gioscia-Ryan R A, Heaton C J, Seals D R (2013). The autophagy enhancer spermidine reverses arterial aging. *Mech Ageing Dev*, 134(7-8):314-320.
- [330] Soda K, Dobashi Y, Kano Y, Tsujinaka S, Konishi F (2009). Polyamine-rich food decreases age-associated pathology and mortality in aged mice. *Exp Gerontol*, 44(11):727-732.
- [331] Yue F, Li W, Zou J, Jiang X, Xu G, Huang H, et al. (2017). Spermidine Prolongs Lifespan and Prevents Liver Fibrosis and Hepatocellular Carcinoma by Activating MAP1S-Mediated Autophagy. *Cancer Res*, 77(11):2938-2951.
- [332] Yuan X, Tian G G, Pei X, Hu X, Wu J (2021). Spermidine induces cytoprotective autophagy of female germline stem cells in vitro and ameliorates aging caused by oxidative stress through upregulated sequestosome-1/p62 expression. *Cell Biosci*, 11(1):107.
- [333] Kim G, Kim M, Kim M, Park C, Yoon Y, Lim D H, et al. (2021). Spermidine-induced recovery of human dermal structure and barrier function by skin microbiome. *Commun Biol*, 4(1):231.
- [334] Wirth M, Benson G, Schwarz C, Kobe T, Grittner U, Schmitz D, et al. (2018). The effect of spermidine on memory performance in older adults at risk for dementia: A randomized controlled trial. *Cortex*, 109(181)-188.
- [335] Binh P, Soda K, Maruyama C, Kawakami M (2010). Relationship between food polyamines and gross domestic product in association with longevity in Asian countries. *Health-London*,
- [336] Yang L, Zhao J, Zhu Y, Tian Z, Wang C (2011). Reactive oxygen species (ROS) accumulation induced by mononaphthalimide-spermidine leads to intrinsic and AIF-mediated apoptosis in HeLa cells. *Oncol Rep*, 25(4):1099-1107.
- [337] Yang L, Li W, Tian Z, Zhao J, Wang C (2011). Mononaphthalimide spermidine conjugate induces cell proliferation inhibition and apoptosis in HeLa cells. *Toxicol in Vitro*, 25(4):882-889.
- [338] Chen Y, Zhuang H, Chen X, Shi Z, Wang X (2018). Spermidine-induced growth inhibition and apoptosis via autophagic activation in cervical cancer. *Oncol Rep*, 39(6):2845-2854.
- [339] Razvi S S, Choudhry H, Moselhy S S, Kumosani T A, Hasan M N, Zamzami M A, et al. (2017). Synthesis, screening and pro-apoptotic activity of novel acyl spermidine derivatives on human cancer cell lines. *Biomed Pharmacother*, 93(190)-201.
- [340] Ohkubo S, Mancinelli R, Miglietta S, Cona A, Angelini R, Canettieri G, et al. (2019). Maize polyamine oxidase in the presence of spermine/spermidine induces the apoptosis of LoVo human colon adenocarcinoma cells. *Int J Oncol*, 54(6):2080-2094.
- [341] Miao H, Ou J, Peng Y, Zhang X, Chen Y, Hao L, et al. (2016). Macrophage ABHD5 promotes colorectal cancer growth by suppressing spermidine production by SRM. *Nat Commun*, 7(1)716.
- [342] Soda K, Kano Y, Chiba F, Koizumi K, Miyaki Y (2013). Increased polyamine intake inhibits age-associated alteration in global DNA methylation and 1,2-dimethylhydrazine-induced tumorigenesis. *Plos One*, 8(5):e64357.

- [343] Vargas A J, Ashbeck E L, Wertheim B C, Wallace R B, Neuhouser M L, Thomson C A, et al. (2015). Dietary polyamine intake and colorectal cancer risk in postmenopausal women. *Am J Clin Nutr*, 102(2):411-419.
- [344] Zagryazhskaya A, Zhivotovsky B (2014). miRNAs in lung cancer: a link to aging. *Ageing Res Rev*, 17(54-67).
- [345] 345. Sorensen SA, Fenger K, Olsen JH (1999). Significantly lower incidence of cancer among patients with Huntington disease: An apoptotic effect of an expanded polyglutamine tract? *Cancer-Am Cancer Soc*, 86(7):1342-1346.
- [346] Musicco M, Adorni F, Di Santo S, Prinelli F, Pettenati C, Caltagirone C, et al. (2013). Inverse occurrence of cancer and Alzheimer disease: a population-based incidence study. *Neurology*, 81(4):322-328.
- [347] Ording A G, Horváth Puhó E, Veres K, Glymour M M, Rørth M, Sørensen H T, et al. (2020). Cancer and risk of Alzheimer's disease: Small association in a nationwide cohort study. *Alzheimer's & Dementia*, 16(7):953-964.
- [348] Freedman D M, Wu J, Daugherty S E, Kuncel R W, Enewold L R, Pfeiffer R M (2014). The risk of amyotrophic lateral sclerosis after cancer in U.S. elderly adults: a population-based prospective study. *Int J Cancer*, 135(7):1745-1750.
- [349] Liu R, Gao X, Lu Y, Chen H (2011). Meta-analysis of the relationship between Parkinson disease and melanoma. *Neurology*, 76(23):2002-2009.
- [350] Fan J, Feng Z, Chen N (2020). Spermidine as a target for cancer therapy. *Pharmacol Res*, 159(104943).
- [351] Bitto A, Ito T K, Pineda V V, LeTexier N J, Huang H Z, Sutlief E, et al. (2016). Transient rapamycin treatment can increase lifespan and healthspan in middle-aged mice. *Elife*, 5(
- [352] Anisimov V N, Zabezhinski M A, Popovich I G, Piskunova T S, Semenchenko A V, Tyndyk M L, et al. (2011). Rapamycin increases lifespan and inhibits spontaneous tumorigenesis in inbred female mice. *Cell Cycle*, 10(24):4230-4236.
- [353] Komarova E A, Antoch M P, Novototskaya L R, Chernova O B, Paszkiewicz G, Leontieva O V, et al. (2012). Rapamycin extends lifespan and delays tumorigenesis in heterozygous p53^{+/-} mice. *Aging (Albany NY)*, 4(10):709-714.
- [354] Popovich I G, Anisimov V N, Zabezhinski M A, Semenchenko A V, Tyndyk M L, Yurova M N, et al. (2014). Lifespan extension and cancer prevention in HER-2/neu transgenic mice treated with low intermittent doses of rapamycin. *Cancer Biol Ther*, 15(5):586-592.
- [355] Khodabakhshi A, Akbari M E, Mirzaei H R, Mehrad-Majd H, Kalamian M, Davoodi S H (2020). Feasibility, Safety, and Beneficial Effects of MCT-Based Ketogenic Diet for Breast Cancer Treatment: A Randomized Controlled Trial Study. *Nutr Cancer*, 72(4):627-634.